



Received: 22 October 2015
Accepted: 30 November 2015
Published: 11 January 2016

*Corresponding author: Piyush Kumar Bhandari, Department of Mathematics, Shrinathji Institute of Technology & Engineering, Nathdwara, Rajasthan 313301, India
E-mail: bhandari1piyush@gmail.com

Reviewing editor:
Regina S. Burachik, University of South Australia, Australia

Additional information is available at the end of the article

APPLIED & INTERDISCIPLINARY MATHEMATICS | RESEARCH ARTICLE

On some inequalities involving Turán-type inequalities

Piyush Kumar Bhandari^{1*} and S.K. Bissu^{2§}

Abstract: Using a new form of the Cauchy–Bunyakovsky–Schwarz inequality, we prove inequalities involving Turán-type inequalities for some special functions.

Subjects: Applied mathematics; Mathematics & statistics; Science

Keywords: a form of Cauchy–Bunyakovsky–Schwarz inequality; Turán-type inequalities; polygamma functions; exponential integral function; Abramowitz’s function

2010 Mathematics subject classifications: Primary 26D07; Secondary 33B15

1. Introduction

The integral representation of well-known Cauchy–Bunyakovsky–Schwarz inequality (see, for instance, Mitrinović, Pečarić, & Fink, 1993) in the space of continuous real-valued functions $C([a, b], \mathbb{R})$ is given by:

$$\left(\int_a^b u^{\frac{1}{2}}(t)v^{\frac{1}{2}}(t) dt \right)^2 \leq \left(\int_a^b u(t) dt \right) \left(\int_a^b v(t) dt \right) \quad (1)$$

It is well known that the Cauchy–Bunyakovsky–Schwarz inequality plays an important role in different branches of modern mathematics such as Hilbert space theory, classical real and complex analysis, numerical analysis, qualitative theory of differential equations and probability and statistics. To date, a large number of generalisations and refinements of this inequality have been investigated in the literature, e.g. (Alzer, 1999; Callebaut, 1965; Masjed-Jamei, 2009; Masjed-Jamei, Dragomir, & Srivastava, 2009; Steiger, 1969; Zheng, 1998).

ABOUT THE AUTHORS

Piyush Kumar Bhandari is working as an assistant professor in the department of Mathematics, Shrinathji Institute of Technology & Engineering, Nathdwara, Rajasthan, India. He received his MSc degree in Mathematics from M.L. Sukhadia University, Udaipur in 2000, cleared CSIR-UGC NET in June 2001. He is pursuing his PhD in the field of “Inequalities and Special Function”.

S.K. Bissu is working as an associate professor in department of Mathematics, Government College, Ajmer. He received his MSc degree in 1987 from University of Rajasthan, Jaipur and PhD degree in 1992 from M.L. Sukhadia University, Udaipur. He has published 18 research papers in national and international journals and has written several books of Board of Secondary Education, Rajasthan, for secondary and senior secondary levels. His area of interest is “Inequalities and Special Function, Fractional calculus”.

PUBLIC INTEREST STATEMENT

In this paper, we prove inequalities involving Turán-type inequalities for some special functions using a new form of the Cauchy–Bunyakovsky–Schwarz inequality. These inequalities play an important role in different branches of modern mathematics such as Hilbert space theory, classical real and complex analysis, numerical analysis, probability and statistics. Also, Turán-type inequalities have important applications in complex analysis, number theory, theory of mean values or statistics and control theory.

Also, the importance, in many fields of mathematics, of the inequalities of the type:

$$f_n(x)f_{n+2}(x) - f_{n+1}^2(x) \leq 0 \tag{2}$$

$n = 0, 1, 2, \dots$ is well known. They are named, by Karlin and Szegő, Turán-type inequalities because the first of this type of inequalities was proved by Turán, (1950).

Laforgia and Natalini, (2006) used the following form of the Schwarz inequality (1):

$$\left(\int_a^b g(t) f^{\frac{m+n}{2}}(t) dt \right)^2 \leq \left(\int_a^b g(t) f^m(t) dt \right) \left(\int_a^b g(t) f^n(t) dt \right) \tag{3}$$

to establish some new Turán-type inequalities involving the special functions as gamma, polygamma functions and Riemann’s zeta function. Here, f and g are non-negative functions of a real variable and m and n belong to a set S of real numbers, such that the involved integrals in Equation (3) exist.

In this context, we have the idea to replace $u(t)$ and $v(t)$ in (1) by $g(t)h^{\alpha x}(t)f^{\nu}(t)$ and $g(t)h^{(2-\alpha)x}(t)f^{\mu}(t)$, respectively, to introduce the following new inequality:

$$\left(\int_a^b g(t)h^{\alpha x}(t)f^{\frac{\nu+\mu}{2}}(t) dt \right)^2 \leq \left(\int_a^b g(t)h^{\alpha x}(t)f^{\nu}(t) dt \right) \left(\int_a^b g(t)h^{(2-\alpha)x}(t)f^{\mu}(t) dt \right) \tag{4}$$

in which $\alpha, \nu, \mu \in \mathbb{R}$ and g, h, f are real integrable functions, such that the involved integrals in Equation (4) exist.

For $h(t) = 1$, or $x = 0$, our new inequality Equation (4) reduces to the inequality Equation (3).

The aim of this paper is to apply the inequality (4) for some well-known special functions in order to get inequalities involving Turán-type inequalities.

2. The results

In this section, we apply the inequality Equation (4) to prove inequalities involving Turán-type inequalities for n -th derivative of gamma function and the Remainder of the Binet’s first formula for $\ln \Gamma(x)$, polygamma functions, exponential integral function, Abramowitz’s function and modified Bessel function of second kind.

2.1. An inequality for the n -th derivative of gamma function

THEOREM 2.1 For every real number $x \in (0, \infty)$, $\alpha \in (0, 2)$ and for every integer $\nu, \mu \geq 1$, such that $\nu + \mu$ is even, it holds for the n -th derivative of gamma function:

$$\left(\Gamma^{(\frac{\nu+\mu}{2})}(x) \right)^2 \leq \Gamma^{(\nu)}(\alpha x) \Gamma^{(\mu)}((2-\alpha)x)$$

Proof The classical Euler gamma function is defined for $x > 0$ as:

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt \tag{5}$$

By differentiating Equation (5), we obtain, for $n = 1, 2, 3, \dots$

$$\Gamma^{(n)}(x) = \int_0^\infty e^{-t} t^{x-1} \log^n(t) dt \tag{6}$$

Hence, if we replace $g(t) = e^{-t}t^{-1}$, $h(t) = t$, $f(t) = \log t$ and $[a, b] = [0, \infty)$ in the inequality Equation (4), we get:

$$\begin{aligned} \left(\int_0^\infty t^{-1} e^{-t} t^x \log^{\frac{\nu+\mu}{2}}(t) dt \right)^2 &\leq \left(\int_0^\infty t^{-1} e^{-t} t^{\alpha x} \log^\nu(t) dt \right) \left(\int_0^\infty t^{-1} e^{-t} t^{(2-\alpha)x} \log^\mu(t) dt \right) \\ \Rightarrow \left(\int_0^\infty e^{-t} t^{x-1} \log^{\frac{\nu+\mu}{2}}(t) dt \right)^2 &\leq \left(\int_0^\infty e^{-t} t^{\alpha x-1} \log^\nu(t) dt \right) \left(\int_0^\infty e^{-t} t^{(2-\alpha)x-1} \log^\mu(t) dt \right) \end{aligned}$$

By applying Equation (6) in the above inequality, the following result will eventually be obtained:

$$\left(\Gamma^{\left(\frac{\nu+\mu}{2}\right)}(x) \right)^2 \leq \Gamma^{(\nu)}(\alpha x) \Gamma^{(\mu)}((2-\alpha)x) \tag{7}$$

$\forall \alpha \in (0, 2)$, $x > 0$ and for every integer $\nu, \mu \geq 1$, such that $\nu + \mu$ is even.

In particular, for $\alpha = 1$ and $\mu = \nu + 2$, it obtains the Turán-type inequality for $\nu \in \mathbb{N}$:

$$\left(\Gamma^{(\nu+1)}(x) \right)^2 \leq \Gamma^{(\nu)}(x) \Gamma^{(\nu+2)}(x)$$

For instance, substituting $\alpha = \frac{1}{2}$, $\nu = 4$ and $\mu = 2$ in Equation (7), we get:

$$\left(\Gamma^{(3)}(x) \right)^2 \leq \Gamma^{(4)}\left(\frac{1}{2}x\right) \Gamma^{(2)}\left(\frac{3}{2}x\right) \quad \forall x > 0 \tag{8}$$

2.2. An inequality for the polygamma function

THEOREM 2.2 For every real number $x \in (0, \infty)$, $\alpha \in (0, 2)$ and for every integer $\nu, \mu \geq 1$, such that $\nu + \mu$ is even, it holds for the polygamma functions:

$$\left(\psi^{\left(\frac{\nu+\mu}{2}\right)}(x) \right)^2 \leq \psi^{(\nu)}(\alpha x) \psi^{(\mu)}((2-\alpha)x).$$

Proof As we know, the polygamma functions $\psi^{(n)}(x) = \frac{d^n \psi(x)}{dx^n}$, where $n = 1, 2, 3, \dots$, are defined as the n -th derivative of the Psi function $\psi(x) = \frac{d}{dx} \ln \Gamma(x) = \frac{\Gamma'(x)}{\Gamma(x)}$, ($x > 0$) with the usual notation for the gamma function and has an integral representation (Nikiforov & Uvarov, 1988) as:

$$\psi^{(n)}(x) = (-1)^{n+1} \int_0^\infty \frac{t^n}{1-e^{-t}} e^{-xt} dt \quad (n = 1, 2, \dots; x > 0). \tag{9}$$

Now, if $g(t) = \frac{1}{1-e^{-t}}$, $h(t) = e^{-t}$ and $f(t) = t$ are substituted in inequality Equation (4) for $[a, b] = [0, \infty)$, the following inequality is derived:

$$\left(\int_0^\infty \frac{t^{\frac{\nu+\mu}{2}}}{1-e^{-t}} e^{-xt} dt \right)^2 \leq \left(\int_0^\infty \frac{t^\nu}{1-e^{-t}} e^{-\alpha xt} dt \right) \left(\int_0^\infty \frac{t^\mu}{1-e^{-t}} e^{-(2-\alpha)xt} dt \right)$$

By the definition Equation (9), this is equivalent to:

$$\left(\psi^{(\frac{\nu+\mu}{2})}(x)\right)^2 \leq \psi^{(\nu)}(\alpha x) \psi^{(\mu)}((2-\alpha)x) \tag{10}$$

$\forall \alpha \in (0, 2), x > 0$ and for every integer $\nu, \mu \geq 1$, such that $\nu + \mu$ is even.

In the particular case, for $\alpha = 1$ and $\mu = \nu + 2$, it obtains the Turán-type inequality for $\nu \in \mathbb{N}$:

$$\left(\psi^{(\nu+1)}(x)\right)^2 \leq \psi^{(\nu)}(x) \psi^{(\nu+2)}(x).$$

2.3. An inequality for the n -th derivative of the remainder of the Binet’s first formula for $\ln \Gamma(x)$

THEOREM 2.3 For every real number $x \in (0, \infty), \alpha \in (0, 2)$ and for every integer $\nu, \mu \geq 1$, such that $\nu + \mu$ is even, it holds for the n -th derivative of the remainder of the Binet’s first formula for the logarithm of the gamma function, i.e. $\ln \Gamma(x)$:

$$\theta^{\frac{\nu+\mu}{2}}(x) \leq \theta^\nu(\alpha x) \theta^\mu((2-\alpha)x).$$

Proof Binet’s first formula for $\ln \Gamma(x)$ is given by:

$$\log \Gamma(x) = (x - 1/2) \log x - x + \log \sqrt{2\pi} + \theta(x)$$

For $x > 0$, where the function:

$$\theta(x) = \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \frac{e^{-xt}}{t} dt \tag{11}$$

is known as the remainder of the Binet’s first formula for the logarithm of the gamma function; see (Abramowitz & Stegun, 1965).

By differentiating Equation (11), we obtain, for every positive integer $n \geq 1$.

$$\theta^{(n)}(x) = (-1)^n \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) t^{n-1} e^{-xt} dt \tag{12}$$

Hence, if $g(t) = \frac{1}{t} \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right), h(t) = e^{-t}, f(t) = t$ and $[a, b] = [0, \infty)$, are considered in inequality Equation (4), then we get:

$$\begin{aligned} & \left\{ \int_0^\infty \frac{1}{t} \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) t^{\frac{\nu+\mu}{2}} e^{-xt} dt \right\}^2 \\ & \leq \left\{ \int_0^\infty \frac{1}{t} \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) t^\nu e^{-\alpha xt} dt \right\} \left\{ \int_0^\infty \frac{1}{t} \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) t^\mu e^{-(2-\alpha)xt} dt \right\} \\ & \Rightarrow \left\{ \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) t^{\frac{\nu+\mu}{2} - 1} e^{-xt} dt \right\}^2 \\ & \leq \left\{ \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) t^{\nu-1} e^{-\alpha xt} dt \right\} \left\{ \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) t^{\mu-1} e^{-(2-\alpha)xt} dt \right\} \end{aligned}$$

By Equation (12), this is transformed to:

$$\left(\theta^{\left(\frac{\nu+\mu}{2}\right)}(x)\right)^2 \leq \theta^{(\nu)}(\alpha x)\theta^{(\mu)}((2-\alpha)x) \tag{13}$$

$\forall \alpha \in (0, 2), x > 0$ and for every integer $\nu, \mu \geq 1$, such that $\nu + \mu$ is even.

In particular, for $\alpha = 1$ and $\mu = \nu + 2$, it obtains the Turán-type inequality for $\nu \in \mathbb{N}$

$$\left(\theta^{(\nu+1)}(x)\right)^2 \leq \theta^{(\nu)}(x)\theta^{(\nu+2)}(x)$$

2.4. An inequality for the exponential integral function

THEOREM 2.4 For every real number $x \in (0, \infty)$, $\alpha \in (0, 2)$ and for every integer $\nu, \mu \geq 0$, such that $\nu + \mu$ is even, it holds for the exponential integral function:

$$\left(E_{\frac{\nu+\mu}{2}}(x)\right)^2 \leq E_{\nu}(\alpha x) E_{\mu}((2-\alpha)x)$$

Proof If we consider the exponential integral function [11, p. 228, 5.1.4] with the following integral representation:

$$E_n(x) = \int_1^{\infty} e^{-xt}t^{-n} dt, (n = 0, 1, \dots; x > 0) \tag{14}$$

and then replace $g(t) = 1$, $h(t) = e^{-t}$ and $f(t) = t^{-1}$ for $[a, b] = [1, \infty)$ in inequality Equation (4), we obtain:

$$\left(\int_1^{\infty} e^{-xt}t^{-\frac{\nu+\mu}{2}} dt\right)^2 \leq \left(\int_1^{\infty} e^{-\alpha xt}t^{-\nu} dt\right)\left(\int_1^{\infty} e^{-(2-\alpha)xt}t^{-\mu} dt\right)$$

Using Equation (14), this is in fact equivalent to:

$$\left(E_{\frac{\nu+\mu}{2}}(x)\right)^2 \leq E_{\nu}(\alpha x) E_{\mu}((2-\alpha)x) \tag{15}$$

$\forall \alpha \in (0, 2), x > 0$ and for every integer $\nu, \mu \geq 0$, such that $\nu + \mu$ is even.

In particular, for $\alpha = 1$ and $\mu = \nu + 2$, it obtains the Turán-type inequality for $\nu \in \mathbb{N}$

$$\left(E_{\nu+1}(x)\right)^2 \leq E_{\nu}(x)E_{\nu+2}(x)$$

2.5. An inequality for the Abramowitz's function

THEOREM 2.5 For every real number $x \geq 0$, $\alpha \in [0, 2]$ and for every non-negative integer ν and μ , such that $\nu + \mu$ is even, it holds for the Abramowitz function:

$$\left(f_{\frac{\nu+\mu}{2}}(x)\right)^2 \leq f_{\nu}(\alpha x) f_{\mu}((2-\alpha)x)$$

Proof The Abramowitz's function (Abramowitz & Stegun, 1965) which has been used in many fields of physics, as the theory of the field of particle and radiation transform, is defined as:

$$f_n(x) = \int_0^{\infty} t^n e^{-t^2-xt^{-1}} dt \tag{16}$$

where n is a non-negative integer and $x \geq 0$.

Now, applying inequality Equation (4) for $g(t) = e^{-t^2}$, $h(t) = e^{-t^{-1}}$, $f(t) = t$ and $[a, b] = [0, \infty)$ results in:

$$\left(\int_0^\infty t^{\frac{\nu+\mu}{2}} e^{-t^2-xt^{-1}} dt \right)^2 \leq \left(\int_0^\infty t^\nu e^{-t^2-\alpha xt^{-1}} dt \right) \left(\int_0^\infty t^\mu e^{-t^2-(2-\alpha)xt^{-1}} dt \right)$$

Therefore, according to Equation (16), one can finally arrive at:

$$\left(f_{\frac{\nu+\mu}{2}}(x) \right)^2 \leq f_\nu(\alpha x) f_\mu((2-\alpha)x) \tag{17}$$

$\forall \alpha \in [0, 2]$, $x \geq 0$ and for every non-negative integer ν and μ , such that $\nu + \mu$ is even.

In particular, for $\alpha = 1$ and $\mu = \nu + 2$, it obtains the Turán-type inequality for $\nu \in \mathbb{N}$:

$$\left(f_{\nu+1}(x) \right)^2 \leq f_\nu(x) f_{\nu+2}(x)$$

2.6. An inequality for modified Bessel function of second kind

THEOREM 2.6 For every real number $x \in (0, \infty)$, $\alpha \in (0, 2)$, $\nu > -1/2$ and $\mu > -1/2$, it holds for the modified Bessel function of second kind:

$$K^2\left\{\frac{\nu+\mu}{2}; x\right\} \leq \frac{\Gamma\left\{\nu+\frac{1}{2}\right\}\Gamma\left\{\mu+\frac{1}{2}\right\}}{(\alpha)^\nu(2-\alpha)^\mu\Gamma^2\left\{\frac{\nu+\mu}{2}+\frac{1}{2}\right\}} K\{\nu; \alpha x\} K\{\mu; (2-\alpha)x\}$$

Proof It is known that the modified Bessel function of second kind (Nikiforov & Uvarov, 1988) can be represented by the following relations for $x > 0$ and $\nu > -1/2$:

$$K_\nu(x) = K(\nu; x) = \frac{\sqrt{\pi}(x/2)^\nu}{\Gamma(\nu+1/2)} \int_1^\infty e^{-xt} (t^2-1)^{\nu-1/2} dt \tag{18}$$

By substituting $g(t) = (t^2-1)^{-1/2}$, $h(t) = e^{-t}$ and $f(t) = (t^2-1)$ in inequality Equation (4) for $[a, b] = [1, \infty)$, we obtain:

$$\left(\int_1^\infty e^{-xt} (t^2-1)^{\frac{\nu+\mu}{2}-\frac{1}{2}} dt \right)^2 \leq \left(\int_1^\infty e^{-\alpha xt} (t^2-1)^{(\nu-1/2)} dt \right) \left(\int_1^\infty e^{-(2-\alpha)xt} (t^2-1)^{(\mu-1/2)} dt \right)$$

Corresponding to definition Equation (18), the following result after simplification eventually yields:

$$K^2\left\{\frac{\nu+\mu}{2}; x\right\} \leq \frac{\Gamma\left\{\nu+\frac{1}{2}\right\}\Gamma\left\{\mu+\frac{1}{2}\right\}}{(\alpha)^\nu(2-\alpha)^\mu\Gamma^2\left\{\frac{\nu+\mu}{2}+\frac{1}{2}\right\}} K\{\nu; \alpha x\} K\{\mu; (2-\alpha)x\} \tag{19}$$

provided that $x > 0$, $\alpha \in (0, 2)$, $\nu > -1/2$ and $\mu > -1/2$.

In the particular case for $\alpha = 1$ and $\mu = \nu + 2$, it obtains the Turán-type inequality:

$$K^2\{\nu+1; x\} \leq \frac{\left(\nu+\frac{3}{2}\right)}{\left(\nu+\frac{1}{2}\right)} K\{\nu; x\} K\{\nu+2; x\}, \quad \forall \nu > -1/2 \tag{20}$$

Acknowledgement

The authors appreciate the anonymous referees for their careful corrections to and valuable comments on the original version of this paper.

Funding

The authors received no direct funding for this research.

Author details

Piyush Kumar Bhandari¹

E-mail: bhandari1piyush@gmail.com

ORCID ID: <http://orcid.org/0000-0001-9656-5441>

S.K. Bissu²

E-mail: susilkbissu@gmail.com

ORCID ID: <http://orcid.org/0000-0001-8856-9872>

¹ Department of Mathematics, Shrinathji Institute of Technology & Engineering, Nathdwara, Rajasthan 313301, India.

² Department of Mathematics, Government College of Ajmer, Ajmer, Rajasthan 305001, India.

§ All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

Citation information

Cite this article as: On some inequalities involving Turán-type inequalities, Piyush Kumar Bhandari & S.K. Bissu, *Cogent Mathematics* (2016), 3: 1130678.

References

Abramowitz, M., & Stegun, I. A. (Eds.). (1965). *Handbook of mathematical functions with formulas, graphs and mathematical tables*. New York, NY: Dover Publications.

Alzer, H. (1999). On the Cauchy-Schwarz inequality. *Journal of Mathematical Analysis and Applications*, 234, 6–14.

<http://dx.doi.org/10.1006/jmaa.1998.6252>

Callebaut, D. K. (1965). Generalization of the Cauchy-Schwarz inequality. *Journal of Mathematical Analysis and Applications*, 12, 491–494.

[http://dx.doi.org/10.1016/0022-247X\(65\)90016-8](http://dx.doi.org/10.1016/0022-247X(65)90016-8)

Laforgia, A., & Natalini, P. (2006). Turán-type inequalities for some special functions. *Journal of Inequalities in Pure and Applied Mathematics*, 27, Article no: 32.

Masjed-Jamei, M. (2009). A functional generalization of the Cauchy-Schwarz inequality and some subclasses. *Applied Mathematics Letters*, 22, 1335–1339.

<http://dx.doi.org/10.1016/j.aml.2009.03.001>

Masjed-Jamei, M., Dragomir, S. S., & Srivastava, H. M. (2009). Some generalizations of the Cauchy-Schwarz and the Cauchy-Bunyakovsky inequalities involving four free parameters and their applications. *Mathematical and Computer Modelling*, 49, 1960–1968.

<http://dx.doi.org/10.1016/j.mcm.2008.09.014>

Mitrinović, D. S., Pečarić, J. E., & Fink, A. M. (1993). *Classical and new inequalities in analysis*. Dordrecht: Kluwer Academic. <http://dx.doi.org/10.1007/978-94-017-1043-5>

Nikiforov, A. F., & Uvarov, V. B. (1988). *Special functions of mathematical physics*. Basel: Birkhäuser.

<http://dx.doi.org/10.1007/978-1-4757-1595-8>

Steiger, W. L. (1969). On a generalization of the Cauchy-Schwarz inequality. *The American Mathematical Monthly*, 76, 815–816. <http://dx.doi.org/10.2307/2317882>

Turán, P. (1950). On the zeros of the polynomials of Legendre. *Casopis Pro Pestovani Matematiky*, 75, 113–122.

Zheng, L. (1998). Remark on a refinement of the Cauchy-Schwarz inequality. *Journal of Mathematical Analysis and Applications*, 218, 13–21. <http://dx.doi.org/10.1006/jmaa.1997.5720>



© 2016 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



Cogent Mathematics (ISSN: 2331-1835) is published by Cogent OA, part of Taylor & Francis Group.

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

