



Received: 19 October 2015
Accepted: 05 April 2016
Published: 04 May 2016

*Corresponding author: J. Vernold Vivin, Department of Mathematics, University College of Engineering Nagercoil (Anna University Constituent College), Konam, Nagercoil 629 004, Tamil Nadu, India
E-mail: vernoldvivin@yahoo.in

Reviewing editor:
Feng Qi, Tianjin Polytechnic University, China

Additional information is available at the end of the article

APPLIED & INTERDISCIPLINARY MATHEMATICS | SHORT COMMUNICATION

δ -Dynamic chromatic number of Helm graph families

N. Mohanapriya^{1,2}, J. Vernold Vivin^{3*} and M. Venkatachalam¹

Abstract: An r -dynamic coloring of a graph G is a proper coloring c of the vertices such that $|c(N(v))| \geq \min\{r, d(v)\}$, for each $v \in V(G)$, where $N(v)$ and $d(v)$ denote the neighborhood and the degree of v , respectively. The r -dynamic chromatic number of a graph G is the minimum k such that G has an r -dynamic coloring with k colors. In this paper, we obtain the δ -dynamic chromatic number of middle, total, and central of helm graph, where $\delta = \min_{v \in V(G)} \{d(v)\}$.

Subjects: Advanced Mathematics; Combinatorics; Discrete Mathematics; Mathematics & Statistics; Science

Keywords: r -dynamic coloring; middle graph; total graph; central graph

2000 Mathematics subject classifications: 05C15

1. Introduction

Throughout this paper all graphs are finite and simple. The r -dynamic chromatic number was first introduced by Montgomery (2001). An r -dynamic coloring of a graph G is a map c from $V(G)$ to a set of colors such that (i) if $uv \in E(G)$, then $c(u) \neq c(v)$, and (ii) for each vertex

ABOUT THE AUTHORS



J. Vernold Vivin

N. Mohanapriya is working as an assistant professor in the Department of Mathematics, RVS Technical Campus—Coimbatore, Tamil Nadu, India. She is pursuing her Part-time (Category B) PhD in Bharathiar University, Coimbatore in the field of graph theory, particularly in graph colorings. She has published five research papers in International and National Journals and Conferences.

J. Vernold Vivin is working as an assistant professor in the Department of Mathematics, University College of Engineering Nagercoil, (Anna University, Constituent College), Konam, Nagercoil. His research areas of interests are graph theory and fractional differential equations. He has published more than 50 research papers in International and National Journals and Conferences.

M. Venkatachalam is working as an assistant professor in the Department of Mathematics, RVS Technical Campus—Coimbatore, Tamil Nadu, India. His research areas of interests are graph theory and inventory models. He has published more than 20 research papers in International and National Journals and Conferences.

PUBLIC INTEREST STATEMENT

Graph coloring is one of the research areas that shaped the graph theory as we know it today; the attempts to prove many more theorems in graph colorings have inspired many notions that became important on their own. Also motivated the study of colorings of various families of graphs, including the graphs embedded in the surfaces of bounded genus. Even though a computer-assisted proof of the theorems was eventually found, many natural problems motivated by it remain unsolved and the study of colorings of planar graphs and of graphs on surfaces is one of the most active areas of research in modern graph theory. This research paper outlines some of recently developed coloring, i.e. dynamic coloring. The new results in this paper are demonstrated by giving detailed proof based on our recent papers and to the best of our knowledge.

$v \in V(G)$, $|c(N(v))| \geq \min\{r, d(v)\}$, where $N(v)$ denotes the set of vertices adjacent to v and $d(v)$ is its degree. The r -dynamic chromatic number of a graph G , written $\chi_r(G)$, is the minimum k such that G has an r -dynamic proper k -coloring. The 1-dynamic chromatic number of a graph G is equal to its chromatic number. The 2-dynamic chromatic number of a graph has been studied under the name dynamic chromatic number in Ahadi, Akbari, Dehghana, and Ghanbari (2012), Akbari, Ghanbari, and Jahanbakam (2009, 2010), Alishahi (2012), Lai, Montgomery, and Poon (2003). There are many upper bounds and lower bounds for $\chi_d(G)$ in terms of graph parameters. For example,

For a graph G with $\Delta(G) \geq 3$, Lai et al. (2003) proved that $\chi_d(G) \leq \Delta(G) + 1$. An upper bound for the dynamic chromatic number of a d -regular graph G in terms of $\chi(G)$ and the independence number of G , $\alpha(G)$, was introduced in Dehghan and Ahadi (2012). In fact, it was proved that $\chi_d(G) \leq \chi(G) + 2 \log_2 \alpha(G) + 3$. Taherkhani gave in (2016) an upper bound for $\chi_2(G)$ in terms of the chromatic number, the maximum degree Δ and the minimum degree δ , i.e. $\chi_2(G) - \chi(G) \leq \lceil (\Delta e) / \delta \log(2e(\Delta^2 + 1)) \rceil$.

Li, Yao, Zhou, and Broersma proved in (2009) that the computational complexity of $\chi_d(G)$ for a 3-regular graph is an NP-complete problem. Furthermore, Liu and Zhou (2008) showed that to determine whether there exists a 3-dynamic coloring, for a claw free graph with maximum degree 3, is NP-complete.

In this paper, we study $\chi_r(G)$ when r is δ , the minimum degree of the graph. We find the δ -dynamic chromatic number for middle, total, and central graph of helm graph.

2. Results

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. The middle graph (Michalak, 1981) of G , denoted by $M(G)$ is defined as follows. The vertex set of $M(G)$ is $V(G) \cup E(G)$. Two vertices x, y of $M(G)$ are adjacent in $M(G)$ in case one of the following holds: (i) x, y are in $E(G)$ and x, y are adjacent in G . (ii) x is in $V(G)$, y is in $E(G)$, and x, y are incident in G .

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. The total graph (Michalak, 1981) of G , denoted by $T(G)$ is defined in the following way. The vertex set of $T(G)$ is $V(G) \cup E(G)$. Two vertices x, y of $T(G)$ are adjacent in $T(G)$ in case one of the following holds: (i) x, y are in $V(G)$ and x is adjacent to y in G . (ii) x, y are in $E(G)$ and x, y are adjacent in G . (iii) x is in $V(G)$, y is in $E(G)$ and x, y are incident in G .

The central graph (Vernold Vivin, 2007) $C(G)$ of a graph G is obtained from G by subdividing each edge of G exactly once and then joining each pair of vertices of the original graph which were previously non-adjacent.

The helm graph H_n is the graph obtained from an n -wheel graph by adjoining a pendent edge at each node of the cycle. Where $V(H_n) = \{v\} \cup \{v_1, v_2, \dots, v_{n-1}\} \cup \{u_1, u_2, \dots, u_{n-1}\}$ and $E(H_n) = \{e_i: 1 \leq i \leq n-1\} \cup \{e'_i: 1 \leq i \leq n-1\} \cup \{s_i: 1 \leq i \leq n-1\}$, where e_i is the edge $v_i v_{i+1}$ ($1 \leq i \leq n-1$), e'_i is the edge $v_i v_{i+1}$ ($1 \leq i \leq n-2$) and s_i is the edge $v_i u_i$ ($1 \leq i \leq n-1$). This notation is valid through the entire paper.

THEOREM 2.1 *Let $n \geq 8$. The δ -dynamic chromatic number of the middle graph of a helm of order $2n - 1$ is $\chi_\delta(M(H_n)) = n$.*

Proof By the definition of middle graph, $V(M(H_n)) = V(H_n) \cup E(H_n) = \{v\} \cup \{v_i: 1 \leq i \leq n-1\} \cup \{u_i: 1 \leq i \leq n-1\} \cup \{e_i: 1 \leq i \leq n-1\} \cup \{e'_i: 1 \leq i \leq n-1\} \cup \{s_i: 1 \leq i \leq n-1\}$.

The vertices v and $\{e_i: 1 \leq i \leq n-1\}$ induce a clique of order K_n in $M(H_n)$. Thus, $\chi_\delta(M(H_n)) \geq n$.

Consider the following n -coloring of $M(H_n)$:

For $1 \leq i \leq n - 1$, assign the color c_i to e_i and assign the color c_n to v . For $1 \leq i \leq n - 1$, assign the color c_n to u_i , $\deg(u_i) = \delta(M(H_n)) = 1$. For $1 \leq i \leq n - 1$, assign to e'_i one of the allowed colors—such color exists, because $\deg(u_i) = 8$. For $1 \leq i \leq n - 1$, if any, assign to vertex v_i one of the allowed colors—such color exists, because $\deg(v_i) = 4$. For $1 \leq i \leq n - 1$, if any, assign to vertex s_i one of the allowed colors—such color exists, because $\deg(s_i) = 3$. An easy check shows that $N(v)$ contains an induced clique of order 5, for every $v \in V(M(H_n))$. Thus, this coloring is a δ -dynamic coloring. Hence, $\chi_\delta(M(H_n)) \leq n$. Therefore, $\chi_\delta(M(H_n)) = n, \forall n \geq 8$. \square

THEOREM 2.2 Let $n \geq 9$. The δ -dynamic chromatic number of the total graph of a helm of order $2n - 1$ is $\chi_\delta(T(H_n)) = n$.

Proof By the definition of total graph $V(T(H_n)) = V(H_n) \cup E(H_n) = \{v\} \cup \{v_i; 1 \leq i \leq n - 1\} \cup \{u_i; 1 \leq i \leq n - 1\} \cup \{e_i; 1 \leq i \leq n - 1\} \cup \{e'_i; 1 \leq i \leq n - 1\} \cup \{s_i; 1 \leq i \leq n - 1\}$. The vertices v and $\{e_i; 1 \leq i \leq n - 1\}$ induce a clique of order K_n in $T(H_n)$. Thus, $\chi_\delta(T(H_n)) \geq n$.

Consider the following n -coloring of $T(H_n)$:

For $1 \leq i \leq n - 1$, assign the color c_i to e_i and assign the color c_n to v . For $1 \leq i \leq n - 1$, assign to u_i one of the allowed colors—such color exists, because $\delta(u_i) = \deg(u_i) = 2$. For $1 \leq i \leq n - 1$, assign to e'_i one of the allowed colors—such color exists, because $\deg(u_i) = 8$. For $1 \leq i \leq n - 1$, if any, assign to vertex v_i one of the allowed colors—such color exists, because $\deg(v_i) = 8$. For $1 \leq i \leq n - 1$, if any, assign to vertex s_i one of the allowed colors—such color exists, because $\deg(s_i) = 5$. An easy check shows that $N(v)$ contains an induced clique of order 5, for every $v \in V(M(H_n))$. Thus, this coloring is a δ -dynamic coloring. Hence, $\chi_\delta(T(H_n)) \leq n$. Therefore, $\chi_\delta(T(H_n)) = n, \forall n \geq 9$. \square

THEOREM 2.3 Let $n \geq 4$. The δ -dynamic chromatic number of the central graph of a helm of order $2n - 1$ is $\chi_\delta(C(H_n)) = 2n - 1$.

Proof By the definition of central graph, subdividing each edge of H_n exactly once and then joining each pair of vertices of H_n which were non-adjacent. Let $V(C(H_n)) = V(H_n) \cup E(H_n) = \{v\} \cup \{v_i; 1 \leq i \leq n - 1\} \cup \{u_i; 1 \leq i \leq n - 1\} \cup \{e_i; 1 \leq i \leq n - 1\} \cup \{e'_i; 1 \leq i \leq n - 1\} \cup \{s_i; 1 \leq i \leq n - 1\}$. Clearly, the graph induced by $\{v_{2i}; i = 1, 2, \dots, \lfloor (n - 1)/2 \rfloor\}$ is a complete graph. Thus, a proper coloring assigns at least $\lfloor (n - 1)/2 \rfloor$ colors to them. The same happens with the subgraph induced by $\{v_{2i-1}; i = 1, 2, \dots, \lfloor (n - 1)/2 \rfloor\}$. Moreover, if we are considering a δ -dynamic coloring when $n - 1$ is odd v_{n-1} should have a different color from $v_{2i-1}, i = 1, 2, \dots, (n - 1)/2$, because v_{n-1} and v_1 are the only neighbors of e'_{n-1} and v_{n-1} is adjacent to $v_{2i-1}, i = 2, \dots, (n - 1)/2$. A similar reasoning also shows that in a δ -dynamic coloring, the colors assigned to odd vertices should be different to the colors assigned to even vertices and that all of them should be different from the color assigned to v . It is also shown that in a δ -dynamic coloring, the colors assigned to vertices $\{v\} \cup \{v_i; 1 \leq i \leq n - 1\}$ should be different to the colors assigned to the vertices of $\{u_i; 1 \leq i \leq n - 1\}$. Since, the vertices $\{u_i; 1 \leq i \leq n - 1\}$ and v induces a clique of order K_n in $C(H_n)$ and the vertices $\{v_i; 1 \leq i \leq n - 1\}$ adjacent to $\{u_j; 1 \leq j \leq n - 1\} \forall i \neq j$. But, any three consecutive vertices of the path must be colored differently in any dynamic coloring. Since, the first and third vertices are the only neighbors of the second vertex and must be colored differently (by the condition of dynamic coloring) and also differently from the second vertex. So, the same color to $\{v_i; 1 \leq i \leq n - 1\}$ and $\{u_j; 1 \leq j \leq n - 1\} \forall i = j$ is impossible. Thus, $\chi_\delta(C(H_n)) \geq 2n - 1$.

Consider the following $2n - 1$ -coloring of $C(H_n)$:

For $1 \leq i \leq n - 1$, assign the color c_i to v_i and assign the color c_n to v . For $1 \leq i \leq n - 1$, assign the color c_{n+i} to u_i . For $1 \leq i \leq n - 1$, assign to vertices s_i, e'_i and e_i one of the allowed colors—such color exists, because $\deg(s_i) = \deg(e'_i) = \deg(e_i) = \delta(C(H_n)) = 2$. An easy check shows that this coloring is a δ -dynamic coloring. Hence, $\chi_\delta(C(H_n)) \leq 2n - 1$. Therefore, $\chi_\delta(C(H_n)) = 2n - 1$. \square

Acknowledgements

With due Respect, the authors sincerely thank the referee for his careful reading, excellent comments and fruitful suggestions that have resulted in the improvement of the quality of this manuscript.

Funding

The authors received no direct funding for this research.

Author details

N. Mohanapriya^{1,2}

E-mail: n.mohanamaths@gmail.com

J. Vernold Vivin³

E-mail: vernoldvivin@yahoo.in

ORCID ID: <http://orcid.org/0000-0002-3027-2010>

M. Venkatachalam¹

E-mail: venkatmaths@gmail.com

¹ RVS Faculty of Engineering, Department of Mathematics, RVS Technical Campus - Coimbatore, Coimbatore 641 402, Tamil Nadu, India.

² Bharathiar University, Coimbatore 641 046, Tamil Nadu, India.

³ Department of Mathematics, University College of Engineering Nagercoil (Anna University Constituent College), Konam, Nagercoil 629 004, Tamil Nadu, India.

Citation information

Cite this article as: δ -Dynamic chromatic number of Helm graph families, N. Mohanapriya, J. Vernold Vivin & M. Venkatachalam, *Cogent Mathematics* (2016), 3: 1178411.

References

Ahadi, A., Akbari, S., Dehghana, A., & Ghanbari, M. (2012). On the difference between chromatic number and dynamic

chromatic number of graphs. *Discrete Mathematics*, 312, 2579–2583.

Akbari, S., Ghanbari, M., & Jahanbekam, S. (2009). On the list dynamic coloring of graphs. *Discrete Applied Mathematics*, 157, 3005–3007.

Akbari, S., Ghanbari, M., & Jahanbekam, S. (2010). On the dynamic chromatic number of graphs. *Combinatorics and Graphs, in: Contemporary Mathematics - American Mathematical Society*, 531, 11–18.

Alishahi, M. (2012). Dynamic chromatic number of regular graphs. *Discrete Applied Mathematics*, 160, 2098–2103.

Dehghan, A., & Ahadi, A. (2012). Upper bounds for the 2-hued chromatic number of graphs in terms of the independence number. *Discrete Applied Mathematics*, 160, 2142–2146.

Lai, H. J., Montgomery, B., & Poon, H. (2003). Upper bounds of dynamic chromatic number. *Ars Combinatoria*, 68, 193–201.

Li, X., Yao, X., Zhou, W., & Broersma, H. (2009). Complexity of conditional colorability of graphs. *Applied Mathematics Letters*, 22, 320–324.

Li, X., & Zhou, W. (2008). The 2nd-order conditional 3-coloring of claw-free graphs. *Theoretical Computer Science*, 396, 151–157.

Michalak, D. (1981). On middle and total graphs with coarseness number equal 1, Springer Verlag graph theory. In *Lagow proceedings* (pp. 139–150). New York: Springer Verlag Berlin Heidelberg.

Montgomery, B. (2001). *Dynamic coloring of graphs* (PhD thesis). West Virginia University. Ann Arbor, MI: ProQuest LLC.

Taherkhani, A. (2016). On r -Dynamic chromatic number of graphs. *Discrete Applied Mathematics*, 201, 222–227.

Vernold Vivin, J. (2007). *Harmonious coloring of total graphs, n -leaf, central graphs and circumdetic graphs* (PhD thesis). Coimbatore: Bharathiar University.

© 2016 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



Cogent Mathematics (ISSN: 2331-1835) is published by Cogent OA, part of Taylor & Francis Group.

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

