

## **Botanical Insecticide as Simple Extractives for Pest Control**

Wafaa M. Hikal, Rowida S. Baeshen, Hussein A.H. Said-Al Ahl

### **Accepted Manuscript Version**

This is the unedited version of the article as it appeared upon acceptance by the journal. A final edited version of the article in the journal format will be made available soon.

As a service to authors and researchers we publish this version of the accepted manuscript (AM) as soon as possible after acceptance. Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). Please note that during production and pre-press, errors may be discovered which could affect the content.

© 2017 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

**Publisher:** Cogent OA

**Journal:** *Cogent Biology*

**DOI:** <http://doi.org/10.1080/23312025.2017.1404274>



## Botanical Insecticide as Simple Extractives for Pest Control

Wafaa M. Hikal<sup>1,2\*</sup>, Rowida S. Baeshen<sup>1</sup>, Hussein A.H. Said-Al Ahl<sup>3</sup>

<sup>1</sup>Department of Biology, Faculty of Science, University of Tabuk, P.O.Box 741, Tabuk 71491, Saudi Arabia.

<sup>2</sup>Parasitology Laboratory, Water Pollution Researches Department, National Research Center, 33 El-Bohouth St., Dokki, Giza, Egypt, 12622

<sup>3</sup>Medicinal and Aromatic Plants Researches Department, National Research Centre, 33 El-Bohouth St., Dokki, Giza, Egypt, 12622

\*Corresponding Author Email: [wafaahikal@gmail.com](mailto:wafaahikal@gmail.com)

**Abstract:** One of the most important global problems is protecting crops from insects. For the control of insects, synthetic chemicals are continuously used, and their toxicity endangers health of farm operators, animals and food consumers. The negative effects on human health led to a resurgence of interest in botanical insecticides due to their minimal costs and ecological side effects. In this we review the use of plant compounds (essential oils, flavonoids, alkaloids, glycosides, esters and fatty acids) having anti-insect effects and their importance as an alternative to the chemical compounds used in the elimination of insects in different ways, namely repellents, feeding deterrents/antifeedants, toxicants, growth retardants, chemosterilants, and attractants. Botanical insecticides affect only target insects, not destroy beneficial natural enemies and provide residue-free food and safe environment. We, therefore, recommend using botanical insecticides as an integrated insect management program which can greatly reduce the use of synthetic insecticides.

**Subjects:** entomology; insect control; chemistry of medicinal plants; natural control

**Keywords:** natural products; insecticides; active constituents; crop protection; insect

## 1. Introduction

Insects are the most diverse species of animals living on earth and can be found in all habitats. Less than 0.5% of the total number of the known insect species are considered pests, and only a few of these can be a serious menace to people. Insects inflict damage to humans, farm animals and crops. Some insects can constitute a major threat to entire countries or a group of nations. One prominent example is the tsetse fly that puts about 100 million people and 60 million head of cattle at risk in sub-Saharan Africa due to the transmission of trypanosomiasis (Imms, 1964; ICIPE, 1997).

Provision of food has always been a challenge facing mankind. A major cornerstone in this challenge is the competition from insects. Herbivorous insects are said to be responsible for destroying one fifth of the world's total crop production annually. Particularly in the tropics and sub-tropics, where the climate provides a highly favorable environment for a wide range of insects. In the developing countries, the problem of competition from insects is further complicated with a rapid annual increase in the human population. Taking into consideration sudden problems caused by drought in places such as Africa, considerable losses of agricultural products add a serious burden to people's daily life. The introduction of alien insects into new habitats due to the global increase of trade and transport causes another dilemma. Insects inflict their damage on stored products mainly by direct feeding. Some species feed on the endosperm causing loss of weight and quality, while other species feed on the germ, resulting in poor seed germination and less viability. Thus, due to damage done by insects, grains lose value for marketing, consumption or planting. In addition to direct consumption of the product, insect pests contaminate their feeding media through excretion, molting, dead bodies and their own existence in the product, which is not commercially desirable. Damage done by insects encourages infection with bacterial and fungal diseases through transmission of their spores (Malek, & Parveen, 1989; Santos, et al., 1990).

In this review, we will focus on biological control of insects in crop production by using botanical products. We provide an overview of botanical insecticides from the chemical point of view and classify their effects on insects.

## 2. Types of botanical insecticides

### 2.1. Essential oils

Plant secondary natural products are natural chemicals extracted from plants and used as an excellent alternative to synthetic or chemical pesticides (Regnault-Roger, & Philogène, 2008; Sithisut et al., 2011). In addition insecticide resistance to synthetic pesticides, which led to a significant losses of food arising from failure of chemicals in pest and annually caused economic losses of several billion dollars worldwide (Shelton et al., 2002; Elzen, & Hardee, 2003; Pereira et al., 2006), Furthermore, United States Food and Drug Administration (FDA) recognized botanical pesticides (essential oils) as safe than synthetic pesticides which caused increases risk of ozone depletion, neurotoxic, carcinogenic, teratogenic and mutagenic effects in non-targets and cross- and multi-resistance in insects (Regnault-Roger et al., 2012).

Essential oils extracted from aromatic plants have increased considerably as insecticides owing to their popularity with organic growers and environmentally conscious consumers. They have repellent, insecticidal, antifeedants, growth inhibitors, oviposition inhibitors, ovicides and growth-reducing effects on a variety of insects (Lu, 1995; Don-Perdo, 1996; Koshier, & Sedy, 2001; Shelton et al., 2002; Elzen, & Hardee, 2003; Tripathi et al., 2003; Pereira et al., 2006; Sithisut et al., 2011; Regnault-Roger et al., 2012). Essential oils possess interesting larvicidal effects on *Limantria dispar* (Lepi-doptera: Lymantridae, gypsy moth) larvae (Moretti et al., 2000), insecticidal activity, repellent properties against ants, cockroach, bedbugs, head lice, flies and moth and toxic to termites. *Mentha piperita* oil repels ants, flies, lice, moths' and is effective against *Callosobruchus maculatus* and *Tribolium castanum* (Kordali et al., 2005).

*Trachyspermum* sp. oil has larvicidal against *Aedes aegypti* and southern house mosquito, *Culex quinquefasciatus* (Tripathi et al., 2000). Nepetalactone, the active constituent in catnip (*Nepeta cataria*) essential oil is highly effective for repelling mosquitoes, bees and other flying insects. It repels mosquitoes more than DEET. It is particularly effective against *Aedes aegypti* mosquito, a vector for yellow fever virus. Oils of *Zingiber officinale* rhizomes and *Piper cubebaberries* were exhibited insecticidal and antifeeding activities against *T. castaneum* and *S. oryzae* (Chaubey, 2012). Tagetes species oil possesses as anti-insect activity on *Ceratitis capitata* and *Triatoma infestans* (López et al., 2011). *Melaleuca alternifolia* essential oil

possesses the fumigant toxicity against *S. zeamais* (Min et al., 2016). Rosemary, oregano, yarrow, eucalyptus and mint oils used as the safe compounds for surface treating or fumigation in cockroach control. Oregano oil used as a potential repellent against *Supella longipalpa* (Sharififard et al., 2016). Kanat et al. (2003) found that essential oils of many plants have insecticidal effects against the larvae of pine processionary moth, *Thaumetopoea pityocampa*. Also, *Laurus nobilis* essential oil was toxic activities against *Rhyzoperthadominica* and *Tribolium castaneum* (Ben Jemba et al., 2012). *Lavandula hybrida*, *Rosmarinus officinalis* and *Eucalyptus globulus* oils was insecticidal on *Acanthoscelides obtectus* adults (Papachristos et al., 2004). Moreover, essential oil of *Tagetes minuta* has toxicity against *Cochliomyia macellaria* (Diptera: Calliphoridae) and as acaricidal, repellent (Chaaban et al., 2017). Eugenol which the principle compound of the essential oils from basil oil have a strong repellent effect on mosquitoes and linalool also in basil oil has toxic effect to the *Bruchid Zabrotes* sub fasciatus and other storage pests (Chogo, & Crank, 1981; Weaver et al., 1991). Essential oil of *Zingiber zerumbet* has repellent against *Lasioderma serricornis* (Wu et al., 2017). *Juniperus procera* essential oil was exhibited significant repellent against the malarial vector *Anopheles arabiensis* (Karunamoorthi et al., 2014). Terpinen-4-ol; 1,8-cineol; verbenone and camphorin Eucalyptus oil were active against *Acanthoscelides obtectus* adults (Tholl, 2006), antifeedant against biting insects, insecticidal agents and prevent mosquito bite (Fradin, & Day, 2002; Isman, & Machial, 2006; Bakkali, et al., 2008; Batish et al., 2008). Lucia et al. (2007) reported that essential oil from *Eucalyptus globules* is toxic to *Aedes aegypti* larvae. Seyoum et al. (2003) reported that burning of leaves of *Eucalyptus citriodora* used as protection against mosquitoes in Africa. Also, CDC (Center for Disease Control and Prevention, USA) recommended the use of lemon eucalyptus oil (with p-menthane-3,8-diol, PMD, as active ingredient) for protection against West Nile virus that causes neurological disease or even death and is spread by mosquitoes (CDC, 2005).

Toloza et al. (2006) concluded the fumigant toxicity/repellent activity of essential oil from *Eucalyptus cinerea*, *Eucalyptus viminalis* and *Eucalyptus saligna*, against permethrin-resistant humanhead lice. The pesticidal and antifeedant activities of eucalyptus oils has been due to 1,8-cineole, citronellal, citronellol, citronellyl acetate, p-cymene, eucamalol, limonene, linalool,  $\alpha$ -pinene,  $\gamma$ -terpinene,  $\alpha$ -terpineol, alloocimene, and aromadendrene components (Watanabe et al., 1993; Li et al., 1996; Cimanga et al., 2002; Batish et al., 2006; Liu et al., 2008). Eucalyptus oils

rich in cineole have affective against varroa mite, *Varroajacobsoni*-an important parasite of honey bee (Calderone, & Spivak, 1995), *Tetranychus urticae*, *Phytoseiulus persimilis* and *Dermatophagoides pteronyssinus* (Choi et al., 2004; El-Zemity et al., 2006). Eucalyptus essential oil significantly reduced the number of tick bites in humans and concluded that it could be used to reduce tick-borne infections (Chagas et al., 2002; Gardulf et al., 2004). Pujiarti and Fentiyanti (2017) reported that *Eucalyptus deglupta* essential oils have repellent activity against *Culex quinquefasciatus* mosquito. Eugenol, isoeugenol, methyleugenol, methyl isoeugenol, coumarin, coniferyl aldehyde, diniconazole, ethyl cinnamate, and rosmarinic acid against red palm weevil possess as antifeedant activity against adults of red palm weevil (*Rhynchophorus ferrugineus*) (Shukla et al., 2012; Aref et al., 2015; AlJabr et al., 2017).

## 2.2. Alkaloid

Alkaloids are the most important group of natural substances playing an important role in insecticidal (Balandrin et al., 1985; Rattan, 2010). Wachira et al. (2014) concluded that pyridine alkaloids extracted from *Ricinus communis* against the malaria vector *Anopheles gambiae*. Furocoumarin and quinolone alkaloids extracted from *Ruta chalepensis* leaves showed larvicidal and antifeedant activities against the larvae *Spodoptera littoralis* (Emam et al., 2009). Acheuk, and Doumandji-Mitiche (2013) found that alkaloids extract of *Pergularia tomentosa* caused antifeeding and larvicidal effects. Lee (2000) concluded that piperonaline and piperidine alkaloids have mosquito larvicidal activity. Alkaloids from *Arachis hypogaea* extract have larvicidal activity against chikungunya and malarial vectors (Velu et al., 2015).

## 2.3. Flavonoids

Flavonoids could be useful in a pest-management strategy. Flavonoids play an important role in the protection of plants against plant feeding insects' and herbivores (Acheuk, & Doumandji-Mitiche, 2013). Both flavonoids and isoflavonoids protect the plant against insect pests by

influencing their behavior, growth and development (Simmonds, & Stevenson, 2001; Simmonds, 2003). Rutin and quercetin-3-glucoside in *Pinus banksiana* inhibit the development and increase the mortality of *Lymantria dispar* (Gould, & Lister, 2006). Quercetin and rutin glycosides in peanuts caused increased mortality of the tobaccoarmyworm (*Spodoptera litura*) (Mallikarjuna et al., 2004). In rice, three flavone glucosides inhibit digestion in insects and function as deterrent agents in *Nilaparvata lugens* and herbivores (Acheuk, & Doumandji-Mitiche, 2013). Diwan, & Saxena (2010) found that flavinoid glycosides isolated from *Tephrosia purpuria* showed insecticidal property on *Callosobruchus maculatus* grubs. Isoflavonoids and proanthocyanidins are other classes of flavonoids responsible for plant protection against insects. For example, naringenin procyanidin inhibits the development of *Aphis craccivora* and herbivores (Acheuk, & Doumandji-Mitiche, 2013). Kumar et al. (2015) concluded that quercetin/azadirachtin insecticide can be a safe and efficient insecticide which improved the activity of *Euphaedra orientalis* and is non toxic to it, and also is environmentally less harmful as it is easily biodegradable. Goławska et al. (2014) found that two polyphenolic flavonoids (flavanone naringenin and flavonol quercetin) used as insecticides of the pea aphid, *Acyrtosiphon pisum* (Hemiptera: Aphididae). Santos et al. (2016) conclude that *Tagetes erecta* and *Tagetes patula* has phytotoxic compounds (flavonoids) that can promote and expand its use as a natural insecticide. Goławska et al. (2008) suggests that quercetin, kaempferol+RCO-, kaempferol, tricetin, apigenin+RCO-, and apigenin are good for control of the insect pests.

Goławska, & Łukasik (2012) showed the effects of isoflavone genistein and flavone luteolin on the feeding behavior of the pea aphid. Flavonoid glycosides in alfalfa affect feeding behavior of pea aphid (Goławska et al., 2010, 2012). Morimoto et al. (2000) showed that flavonoids can act as feeding deterrents. Flavonoids from *Arachis hypogaea* extract have larvicidal activity against chikungunya and malarial vectors (Velu et al., 2015).

#### **2.4. Glycosides**

Cyanogenic glucosides presented in plant species and considered to have an important role in plant defense against herbivores (Zagrobelny et al., 2004). Al-Rajhy et al. (2003) concluded that the cardiac glycoside, digitoxin, from *Digitalis purpurea*, a cardiac glycosidal (cardenolide)

extract from *Calotropis procera*, azadirachtin and neem oil from *Azadirachta indica* possess against larvae and adult stages of the camel tick, *Hyalomma dromedarii*. Also, Kubo & Kim (1989) found that flavan glycosides, Viscutin-1, -2 and -3, which exhibit insect growth inhibitory activity against the cotton pest insect; *Pectinophora gossypiella*. However, iridoid glycosides possess as insect antifeeding on gypsy moths (*Lymantria dispar*, Lymantriidae) and buckeyes (*Junonia coenia*, Nymphalidae) (Bowers, & Puttick, 1989). Also, cyanogenic glycosides are known as plant defense chemicals and found in cassava, bamboo, flax, and other plants. They are effective against stored-product insects as fumigants. Due to their insecticidal activity to insects, cyanohydrins can be used as an alternative fumigant and also as soil fumigants (Park, & Coats, 2002). Dave & Lediwane (2012) found that anthraquinones isolated from *Cassia* species possess as antimalarial and insecticidal activity. Glycosides from *Arachis hypogaea* extract have larvicidal activity against chikungunya and malarial vectors (Velu et al., 2015). Juvenogens have a potential application in insect pest control (Wimmer et al., 2007).

## 2.5. Esters and fatty acids

Allyl cinnamate caused rapid toxic effects in *Spodoptera littoralis* larvae at low concentrations potential for use in pest control (Giner et al., 2012). Schmidt et al. (2008) demonstrated that ethyl (*E,Z*)-2,4-decadienoate (pear ester) have insecticide against *Cydia pomonella*.

Fatty acids methyl esters were isolated from *Solanum lycocarpum* have larvicidal activity against the vector *Culex quinquefasciatus* (Silva et al., 2015). Mullens et al. (2009) showed that saturated fatty acids (particularly C8, C9 and C10) used as repellents or antifeedants against houseflies, horn flies and stable flies. Samuel et al. (2015) concluded that fatty acids mixture (C8910) has toxicity and repellence against insecticide susceptible and resistant strains of the major malaria vector *Anopheles funestus*. Yousef et al. (2013) reported that toxicity and reduction in larval body weight of linoleic acid against the larvae of *Spodoptera littoralis*.

### 3. Insecticide effects on insects

Botanical insecticides affect various insects in different ways depending on the physiological characteristics of the insect species as well as the type of the insecticidal plant. The components of various botanical insecticidal can be classified into six groups namely; repellents, feeding deterrents/antifeedants, toxicants, growth retardants, chemosterilants, and attractants (Rajashekar et al., 2012).

#### 3. 1. Repellents

A botanical pesticide have a repellent property, where keeps away the insect pest, and protect the crops (Isman, 2006) with minimal impact on the ecosystem, as they drive away the insect pest from the treated materials by stimulating olfactory or other receptors (Talukder et al., 2004; Talukder, 2006). Botanical pesticides are considered safe in pest control because they have low or none pesticide residue making them safe to the people, environment and ecosystem (Talukder et al., 2004). Ghavami et al. (2017) found that essential oils of *Ziziphora tenuiore*, *Myrtus communis*, *Achillea wilhelmsii* and *Mentha piperita* have repellent activities against human fleas. Rahdari, & Hamzei (2017) demonstrated the efficacy of *Mentha piperita*, *Rosmarinus officinalis* and *Coriandrum sativum* oils for applying in organic food protection due to repellent activity of essential oils on *Tribolium confusum*. Zhang et al. (2017) reported the repellent activities of six *Zanthoxylum* species including *Z. armatum*, *Z. dimorphophyllum*, *Z. dimorphophyllum* var. *spinifolium*, *Z. piasezkii*, *Z. stenophyllum*, and *Z. dissitum* essential oils against two storage pests including *T. castaneum* and *L. serricorne* adults and the essential oils of these six *Zanthoxylum* species essential oils possessed significant repellent activities against *T. castaneum* and *L. serricorne* adults. The different repellent activities on two insects might be attributed to the different anti-insect mechanism and different non persistent volatility of essential oil sample. Kimutai et al. (2017) demonstrated that the essential oils of *Cymbopogon citratus* and *Tagetes minuta* on the sandfly, *Phlebotomus duboscqi*. However, the effectiveness of the repellents depends on multiple factors including the type of repellents (active ingredients), formulation, mode of application, environmental factors (temperature, humidity, and wind), the attractiveness of individual people to insects, loss due to removal by perspiration and abrasion, the sensitivity of the insects to repellents, and the biting density. In addition, *Origanum onites*

essential oil had repellent activity against *Amblyomma americanum* (L.) and *Aedes aegypti* (L.). Carvacrol and thymol were strongly repellent to *Aedes aegypti* and *Amblyomma americanum*. Thus, carvacrol-rich *Origanum onites* essential oil, carvacrol and possibly thymol appear to have potential for use to protect humans and domestic animals against mosquitoes and ticks (John et al., 2017). Various natural fatty acids have insecticidal properties, some involving action on acetylcholinesterase and octopaminergic receptors (Perumalsamy et al. 2015). A saturated fatty acid mixture composed of octanoic acid (also called caprylic acid, nonaioic acid and decanoic acid (also called capric acid), collectively called “C8910 acids” (C8, C9, and C10 mixture), repelled horn flies. C8910 acids deterred horn flies from feeding by >85% and the pest was also strongly repelled (Zhu et al. 2015). Feeding deterrence caused by C8910 acids and >50% antifeedancy was observed (Zhu et al. 2015). The naturally occurring oleic and linoleic acids, and methyl oleate synergized the repellency of DEET and the monoterpenoids cuminyl alcohol, cuminaldehyde, and  $\alpha$ -phellandrene (Hieu et al. 2015; Showler, 2017).

### 3. 2. Feeding deterrents/antifeedants

Botanical pesticides that inhibit feeding or disrupt insect feeding by rendering the treated materials unattractive or unpalatable (Rajashekar et al., 2012; Talukder, 2006). The insects remain on the treated material indefinitely and eventually starve to death. Liao et al. (2017) demonstrated that oil of *Melaleuca alternifolia* and their chemical constituents possessed obvious antifeedant activities against *Helicoverpa armigera* Hubner. The phytoconstituents found in the leaf extract of *Khaya senegalensis* include tannins, saponins, flavonoids, steroids, and alkaloids may have been responsible for the mortality of *Dinoderus porcellus* (Loko et al., 2017). Chaudhary et al. (2017) and Ghoneim and Hamadah (2017) pointed that azadirachtin which is prominent constituent of neem established as a pivotal insecticidal ingredient. It acts as an antifeedant, repellent, and repugnant agent and induces sterility in insects by preventing oviposition and interrupting sperm production in males. The same result obtained by Abdullah et al. (2017) who found that, 1,8-cineol found in Galangal essential oil exhibited antifeedant activity, repellent activity, and toxicity effect toward the termites. Jose & Sujatha (2017) revealed that terpenoids, coumarin and phenols present in the methanol extracts of *Gliricidium sepium* exhibited significant antifeedant activity. This indicated that the active compounds

present in the plant inhibit the larval feeding behaviour while others disrupt hormonal balance or make the food unpalatable. These active substances may directly act on the chemosensilla of the larvae resulting in feeding deterrence.

### 3.3. Toxicity

Some botanical pesticides are toxic cause death to stored product insects (Padin et al., 2013). Rotenone is considered as a toxic compound since it is a mitochondrial poison which blocks the electron transport chain and prevents energy production (Hollingworth et al., 1994). As an insecticide, it is a stomach poison because it must be ingested to be effective (Isman, 2006). Essential oil of *Lavandula angustifolia* exhibited good fumigant and contact toxicity against granary weevil adults. In addition, a strong repellent activity able to disrupt granary weevil orientation to an attractive host substrate (Germinara et al., 2017). Trivedi et al. (2017) demonstrated fumigant toxicity against the stored grain pest *Callosobruchus chinensis*. The essential oils of cinnamon, clove, rosemary, bergamot and Japanese mint showed potential to be developed as possible natural fumigants or repellents for control of the pulse beetle. Lucia et al. (2017) found that the mortality of adults and eggs for head lice associated with the use of (geraniol, citronellol, 1,8-cineole, linalool,  $\alpha$ -terpineol, nonyl alcohol, thymol, menthol, carvacrol and eugenol) essential oils. Bouguerra et al. (2017) showed that *Thymus vulgaris* essential oil exhibited significant activity and could be considered as potent natural larvicidal agent against *Cx pipiens*. Zhao et al. (2017) indicated that the essential oil of *Echinops grijsii* roots and the isolated thiophenes have an excellent potential for use in the control of *Aedes albopictus*, *Anopheles Sinensis* and *Culex pipiens pallens* larvae and could be used in the search for new, safer and more effective natural compounds as larvicides. Wu et al. (2017) observed the toxicity and repellent activities of the rhizomes of *Zingiber zerumbet* (L.) Smith (Zingiberaceae) essential oil contains the component  $\alpha$ -caryophyllene against cigarette beetles (*Lasioderma serricornis*). Alkan et al. (2017) indicate that *Heracleum platytaenium* and *Humulus lupulus* extracts have great potentials as insecticides in the management of larvae of *Leptinotarsa decemlineata*. Papanastasiou et al. (2017) showed the toxicity of limonene, linalool and  $\alpha$ -pinene on adult Mediterranean fruit flies. Qari ET AL. (2017) showed DNA damage due to alterations in enzymatic system (acetylcholinesterase, acid phosphatase, alkaline phosphatase, lactate dehydrogenase and phenol oxidase), total protein and DNA concentration after treatment with

essential oils of *Citrus aurantium*, *Eruca sativa*, *Zingiber officinale* and *Origanum majorana* against *Rhyzopertha dominica*. Park et al. (2017) demonstrated that *Thymus vulgaris* oil had the highest insecticidal toxicity followed by *R. graveolens*, *Citrus aurantium*, *L. petersonii* and *A. millefolium* oils. The insecticidal toxicity of *Thymus vulgaris* oil against *P. shantungensis* nymphs was about 1.3-fold more than that against *P. shantungensis* adults. Differences in the insecticidal toxicities of plant-derived oils may be explained on the basis of species-specific responses to plant species, phytochemicals, and the weight and size of *P. shantungensis* adults and nymphs.

### **3. 4. Growth retardants and Development Inhibitors**

Botanical pesticides showed deleterious effects on the growth and development of insects, reducing the weight of larva, pupa and adult stages and lengthening the development stages (Talukder, 2006). Plant derivatives also reduce the survival rates of larvae and pupae as well as adult emergence (Koul et al., 2008). It has been reported that both azadirachtin and neem seed oil increased aphid nymphal mortality significantly at 80% and 77% respectively, and at the same time increasing development time of those surviving to adulthood (Kraiss & Cullen, 2008). Many botanical pesticides have been reported to have a pronounced effect on the developmental period, growth and adult emergence (Shaalán et al., 2005).

### **3.5. Sterility /Reproduction Inhibitors**

Sterility can be induced by sterile insect technique (SIT) or a chemosterilant, a chemical compound that interferes with the reproductive potential of sexually reproducing organism (Morrison et al., 2010). Chemosterilants are used to control economically destructive or disease-causing pests (usually insects) by causing temporary or permanent sterility of one or both of the sexes or preventing maturation of the young to a sexually functional adult stage (Wilke et al., 2009; Navarro-Lopis et al., 2011). It has been reported that plant parts, oil, extracts, and powder mixed with grain reduced insect oviposition, egg hatchability, postembryonic and progeny development (Asawalam & Adesiyán, 2001; Shaalan et al., 2005). Hexane extracts of *Andrographis lineat*, *A. paniculata* and *Tagetes erecta* showed 100% ovicidal activity against *Anopheles subpictus* (Elango et al., 2009). Some botanical insecticides are used as chemosterilants, for example, at the physiological level azadirachtin blocks the synthesis and

release of molting hormones from the prothoracic gland, leading to incomplete ecdysis in immature insects and in adult insects it leads to sterility (Isman, 2006).

Garlic essential oil and its constituents, diallyl sulfide and diallyl disulfide have been highly toxic to *S. zeamais* and *T. castaneum* (Ho et al., 1996; Huang et al., 2000) at different developmental stages. Plata-Rueda et al. (2017) showed that *Tenebrio molitor* was more susceptible in the pupal stage followed by larvae and adults exposed to diallyl sulfide and diallyl disulfide. One possible explanation for the developmental stages difference is that efficacy may be affected by the penetration of the garlic compounds into the body and the ability of the insect to metabolize these compounds. When insects exposed to the garlic essential oil displayed altered locomotion activity, and muscle contractions and paralysis were observed. Paralysis and muscle contractions can be explained by the toxic effect in the nervous system. The toxicity of essential oils in insects indicates neurotoxic action with hyperactivity, hyperextension of the legs and abdomen and rapid knock-down effect or immobilization (Prowse et al., 2006; Zhao et al., 2013). Acetylcholinesterase is an enzyme that has been shown to be inhibited by garlic compounds and can act only or in synergism as diallyl disulfide, diallyl trisulfide, and allicin (Bhatnagar-Thomas et al., 1974; Singh, & Singh, 1996). Diallyl sulfide in garlic compounds have toxic effect in *T. molitor* and may cause inhibition by cross-linking with essential thiol compounds in enzyme structures, altering the functional shape of the protein and denaturalization (Halliwell, & Gutteridge, 1999). Also, diallyl disulfide as main volatile compound in garlic essential oil has repellent properties to *S. zeamais* and *T. castaneum* (Huang et al., 2000). Diallyl disulfide, and diallyl sulfide have high activities of behavioral deterrence against *Tenebrio molitor*, as evaluated by the behavioral responses of larvae and adults to different odor sources and the number of insects repelled, indicating their potential to the pest control in stored products. As well as garlic essential oil compromised the respiration rate of *Tenebrio molitor*. Thus, low respiration rate is an indicator of physiological stress, and essential oils can compromise insect respiration by impairing muscle activity, leading to paralysis (Guedes et al., 2006; Correa et al., 2015; de Araújo et al., 2016).

Dehghani-Samani et al. (2015) showed that essential oil of *Eucalyptus globulus* had repellent activity against *Dermapteran* *gallinae* due to the essential oil components such as 1, 8-cineole, citronellal, citronellol, citronellyl acetate, p-cymene, eucamalol, limonene, linalool,  $\alpha$ -pinene, g-

terpinene,  $\alpha$ -terpineol, alloocimene, and aromadendrene. Among the various components of eucalyptus oil, 1, 8-cineole is the most important one which is largely responsible for a variety of its pesticidal properties and insecticide effects. Jayakumar et al. (2017) recorded the repellent activity of camphor, citronella, eucalyptus, lemon and wintergreen oil essential oils against stored product pests; the adult rice weevil *Sitophilus oryzae* Linnaeus 1763 (Coleoptera: Curculionidae) due to essential oils components repellent nor attractant activity. The protection of stored products and the phytochemical constituents from essential oils can work synergistically, improving their effectiveness.

Ho et al. (1995) concluding that adult mortality might be attributed to the contact toxicity or to the abrasive effect on the pest cuticle (Mathur et al., 1985), which might also interfere with the respiratory mechanism of insect (Schoonhoven, 1978; Agarwal et al., 1988; Kim et al., 2003). Fumigation studies showed that the essential oils had a 'knock down effect' on the test insect. Essential oils act by inhibiting insect acetylcholinesterase (AChE) and thus, ultimately blocking the nerve functions. Also, Obeng-Ofori and Amitaye (2005) who observed signs of immobilization with flexed legs and clinging to the grain outstretched meta thoracic wings from the elytra and paralysis of the dead or dying insects. The enzyme AChE is also the target site of inhibition by organophosphates and carbamate insecticides (Matsumura, 1985). The observed rapid action of essential oils could be attributed to their property of acting in the vapour phase, hence gaining entry into the insect's internal systems with ease through the spiracles; whereas, in topical application procedures, the insect is protected by its exoskeleton against external influences.

Essential oils possess acute contact and fumigant toxicity to insects (Abdelgaleil et al., 2009), repellent activity (Nerio et al., 2009), antifeedant activity (Huang et al., 2000), as well as development and growth inhibitory activity (Waliwitiya et al., 2008). Repellent activity has been linked to the presence of essential oils that cause death of insects by inhibiting AChE activity in the nervous system (Houghton et al., 2006). Essential oils being more useful as insect fumigants (Weaver et al., 1991; Regnault-Roger et al., 1993) and have strong toxicity to insects due to high volatility and lipophilic properties can penetrate into insects rapidly and interfere in physiological functions (Lee et al., 2002; Negahban et al., 2007). Due to their high volatility, they have fumigant and gaseous action on stored product insects. Carvacrol component has broad insecticidal and acaricidal activity against agricultural, storedproduct, medical pests and acts as a

fumigant being highly toxic to adults of *Sitophilus oryzae* (Ahn et al., 1998). Besides, menthol, methonene, limonene,  $\beta$ -pinene,  $\alpha$ -pinene, pulegone, linalool and linalyl acetate exhibited fumigant toxicity in *Sitophilus oryzae* and inhibited AChE activity (Singh et al., 1989; Lee et al., 2001a, b; Koul et al., 2008). Caryophyllene, a volatile compound, was reported to be a strong fumigant and toxic to *Sitophilus zeamais* (Chu et al., 2010).

The insecticidal activity varies with plant derived material, insect species and exposure time. The presence of volatile compounds having strong odour would have blocked the tracheal respiration of the insects leading to their death. Similar observation was made by Liu and Ho (1999) against *Sitophilus zeamais* and *Tribolium castaneum*. Brown (1951) however, pointed out that the amount of fumigant absorbed depends on whether the insect's initial contact with the fumigant resulted in supplication or stimulation of the tracheal opening. Moreover, the ability of the insect to exclude vapor from its cuticle and prevent dehydration of body fluid plays a vital role in susceptibility or tolerance to fumigants of various life stages of insects particularly beetles and weevils infesting stored products (El-Nahal et al., 1989). The toxic effect of essential oils, apart from the variability of phytochemical patterns, involves several other factors. The point of entry of the toxin is one of them where essential oils can be inhaled, ingested or skin absorbed by insects (Regnault-Roger, 1997).

The presence of volatile compounds is responsible for strong odor that could block the tracheal respiration of the insects leading to their death (Pugazhvendan et al., 2012). The mode of action of oils was partially attributed to interference in normal respiration, resulting in suffocation (Schoonhoven, 1978). Most insects breathe through the trachea which usually leads to the opening of the spiracle.

These spiracles might have been blocked thereby leading to suffocation (Adedire et al., 2011; Ileke & Olotuah, 2012) Essential oils are presumed to interfere with basic metabolic, biochemical, physiological and behavioural functions of insects (Mann & Kaufman, 2012). Essential oils block the spiracles, resulting in blockage of respiratory siphons (asphyxiation) and death (Kaufmann & Briegel, 2004; Rotimi et al., 2011). Further, Rattan (2010) reviewed the mechanism of action of essential oils on the body of insects and documented several physiological disruptions, such as inhibition of AChE, disruption of the molecular events of morphogenesis and alteration in the behaviour and memory of cholinergic system. Of these, the

most important activity is the inhibition of AChE activity as it is a key enzyme responsible for terminating the nerve impulse transmission through synaptic pathway. Plant oils affect AChE and have an action on the nervous system (Mikhael, 2011). Recent research has demonstrated the interference of Monoterpenes with AChE activity in insects (Chaubey, 2012a, b). Essential oils are lipophilic in nature and can be inhaled or ingested. The rapid action against insect pests is indicative of a neurotoxic mode of action and interference with the neuromodulator octopamine (Enan, 2005) or GABA-gated chloride channels (Priestley et al., 2006).

Several essential oil components act on the octopaminergic system of insects. Octopamine is a neurotransmitter, neurohormone, and circulating neurohormone-neuromodulator, and its disruption results in total breakdown of the nervous system (Hollingworth et al., 1984). Thus, the octopaminergic system of insects represents a target for insect control. Low molecular weight terpenoids are too lipophilic to be soluble in the haemolymph after crossing the cuticle, and the proposed route of entry is tracheae (Veal, 1996) and may also bind to target sites on receptors that modulate nervous activity (Hollingworth et al., 1984) and interrupt normal neurotransmission leading to paralysis and death.

### **3.6. Attractants**

Botanicals chemicals that cause insects to make oriented movements towards their source are called insect attractants. They influence both gustatory (taste) and olfactory (smell) receptor or sensilla. Iso-thiocyanates from seeds of *Crucifera*, sugar and molasses and terpenes from bark in with pheromones are natural attractants for various insects of *Cruciferaea* and bark beetles. In onion propylmercaptan from *Umbelliferae* and phenylacetaldehyde from flowers of *Araujia serisofera* are attracted carrot fly (*Psila rosae*) and Lepidoptera, respectively. Insect attractants can be used in three ways for the control of insects. In sampling or monitoring insect populations to assess the extent of infestation and decides the measure of control to be adapted in lusting insect to insecticide-coated traps or poison baits and in distracting insects from normal mating, aggregation feeding or oviposition. They do not kill the insects therefore, do not disturb ecosystem. They can be used to misguide the insects to wrong oviposition sites whereby their number will go down by starvation or by producing unfertilized eggs. They cannot be relied as sole control measure used only in integrated control program (Arora et al., 2012).

## **Conclusion**

Botanical insecticides are natural chemicals extracted from plants with insecticidal properties and used as an excellent alternative to synthetic or chemical pesticides for crop protection to avoid negative or side effects of synthetic insecticides. Botanical pesticides (essential oils, flavonoids, alkaloids, glycosides, esters and fatty acids) have various chemical properties and modes of action and affect on insects in different ways namely; repellents, feeding deterrents/antifeedants, toxicants, growth retardants, chemosterilants, and attractants. So it is preferable to use the botanical insecticides instead of synthetic insecticide and these botanical insecticides are recognized by organic crop producers in industrialized countries. So, we recommended using botanical insecticidal and being promoted and research is being conducted to find new sources of botanical insecticides.

## **Conflict of Interests**

The authors declare that there is no conflict of interests regarding the publication of this paper.

## References

- Abdelgaleil, S., Mohamed, M., Badawy, M., & El-arami, S. (2009). Fumigant and contact toxicities of monoterpenes to *Sitophilus oryzae* (L.) and *Tribolium castaneum* (Herbst) and their inhibitory effects on acetylcholinesterase activity. *Journal of Chemical Ecology*, 35:518-525.
- Abdullah, F., Subramanian, P., Ibrahim, H., Abdul Malek, S.N., Lee, G.S., & Hong, S.L. (2017). Chemical composition, antifeedant, repellent, and toxicity activities of the rhizomes of galangal, *Alpinia galanga* Against Asian Subterranean Termites, *Coptotermes gestroi* and *Coptotermes curvignathus* (Isoptera: Rhinotermitidae). *J. Insect Sci.*, 15(7): 2015.
- Acheuk, F., & Doumandji-Mitiche, B. (2013). Insecticidal activity of alkaloids extract of *Pergularia tomentosa* (Asclepiadaceae) against fifth instar larvae of *Locusta migratoria cinerascens* (Fabricius 1781) (Orthoptera: Acrididae). *International Journal of Science and Advanced Technology*, 3, 6: 8-13.
- Adedire, C.O., Obembe, O.M., Akinkurolere, R.O., & Oduleye, S.O. (2011). Response of *Callosobruchus maculatus* Fabricius (Coleoptera: Chrysomelidae: Bruchinae) to extracts of cashew kernels. *Journal of Plant Diseases Protection*, 118, 2:75-79.
- Agarwal, A., Lal, S., & Gupta, K.C. (1988). Natural products as protectants of pulse beetles. *Bulletin of Grain Technology*, 26:154-164.
- Ahn, Y.J., Lee, S.B., Lee, H.S., & Kim, G.H. (1998). Insecticidal and acaricidal activity of carvacrol and b-thujaplicine derived from *Thujopsis dolabrata* var. hondai sawdust. *Journal of Chemical Ecology*, 24:1-90.
- AlJabr, A.M., Hussain, A., Rizwan-ul-Haq, M., & Al-Ayedh, H. (2017). Toxicity of plant secondary metabolites modulating detoxification genes expression for natural red palm weevil pesticide development. *Molecules*, 22, 169.

- Alkan, M., Gokce, A., & Kara, K. (2017). Contact Toxicity of Six Plant Extracts to Different Larval Stages of Colorado Potato Beetle (*Leptinotarsa decemlineata* SAY (Col: Chrysomelidae)). *J. Agr. Sci.*, 23, 309-316.
- Al-Rajhy, D.H., Alahmed, A.M., Hussein, H.I., & Kheir, S.M. (2003). Acaricidal effects of cardiac glycosides, azadirachtin and neem oil against the camel tick, *Hyalomma dromedarii* (Acari: Ixodidae). *Pest Manag. Sci.*, 59, 11:1250-4.
- Aref, S. P., Valizadegan, O., & Farashiani M. E. (2015). *Eucalyptus dundasii* Maiden essential oil, chemical composition and insecticidal values against *Rhyzopertha dominica* (F.) and *Oryzaephilus surinamensis* (L.). *Journal of Plant Protection Research*, 55, 35-41.
- Arora, R., Singh, B., & Dhawan, A.K. (2012). Theory And Practice Of Integrated Pest Management. Scientific Publishers; 5A, New Pali Road, P.O. Box 91, Jodhpur 342 001, India.
- Asawalam, E., & Adesiyan, S. (2001). Potential of *Ocimum basilicum* (Linn) for the control of maize weevil *Sitophilus zeamais* (Motsch). *Nigeria Agricultural Journal*, 32, 1:195-201.
- Bakkali, F., Averbeck, S., Averbeck, D., & Idaomar, M. (2008). Biological effects of essential oils-a review. *Food and Chemical Toxicology*, 46, 446–475.
- Balandrin, M.F., Klocke, J.A., Wurtele, E.S., & Bollinger, W.H. (1985). Natural plant chemicals: sources of industrial and medicinal materials. *Science*, 7; 228(4704):1154-60.
- Batish, D.R., Singh, H.P., Kohli, R.K., & Kaur, S. (2008). Eucalyptus essential oil as a natural pesticide. *Forest Ecology and Management*, 256, 2166–2174.
- Batish, D.R., Singh, H.P., Setia, N., Kaur, S., & Kohli, R.K. (2006). Chemical composition and phytotoxicity of volatile essential oils from intact and fallen leaves of *Eucalyptus citriodora*. *Zeitschrift für Naturforschung C*, 61: 465–471, 2006.
- Ben Jemba, J.M., Tersim, N., Toudert, K.T., & Khouja, M.L. (2012). Insecticidal activities of essential oils from leaves of *Laurus nobilis* L. from Tunisia, Algeria and Morocco, and comparative chemical composition. *Journal of Stored Products Research*, 48, 97–104.

- Bhatnagar-Thomas, P. L. & Pal, A. K. (1974). Studies on the insecticidal activity of garlic oil 2. Mode of action of the oil as a pesticide in *Musca domestica* nebulosa Fabr and *Trogoderma granarium* Everts. *J. Food Sci. Technol.*, 11, 153–158.
- Bouguerra, N., Djebbar, F.T., & Soltani, N. (2017). Algerian *Thymus vulgaris* essential oil: chemical composition and larvicidal activity against the mosquito *Culex pipiens* *International Journal of Mosquito Research*, 4, 1: 37-42.
- Bowers, M. D., & Puttick, G. M. (1989). Iridoid glycosides and insect feeding preferences: gypsy moths (*Lymantria dispar*, Lymantriidae) and buckeyes (*Junonia coenia*, Nymphalidae). *Ecological Entomology*, 14: 247–256.
- Brown AWA. (1951). Insect control by chemicals. New York: Wiley; London: Chapman & Hall, 1951. 817 pp.
- Calderone, N.W., & Spivak, M. (1995). Plant extracts for control of the parasitic mite *Varroa jacobsoni* (Acari: Varroidae) in colonies of the western honey bee (Hymenoptera: Apidae). *Journal of Economic Entomology*, 88, 1211–1215.
- CDC. Center for Disease Control and Prevention, USA (2005). CDC Adopts New Repellent Guidance for Upcoming Mosquito Season. Available at <http://www.cdc.gov/ncidod/dybid/westnile/RepellentUpdates.htm>.
- Chaaban, A., de Souza, A.L.F., Martins, C.E.N., Bertoldi, F.C., & Molento, M.B. (2017). Chemical composition of the essential oil of *Tagetes minuta* and its activity against *Cochliomyia macellaria* (Diptera: Calliphoridae). *European J. Med. Plants*, 18, 1: 1-10.
- Chagas, A.C.S., Passos, W.M., Prates, H.T., Leitem, R.C., Furlong, J., & Fortes, I.C.P. (2002). Acaricide effect of *Eucalyptus* spp. essential oils and concentrated emulsion on *Boophilus microplus*. *Braz. J. Vet. Res. An. Sci.*, 39, 247–253.
- Chaubey, M.K. (2012, a). Responses of *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae) against essential oils and pure compounds. *Herba Polonica*, 58, 3: 33-45.

- Chaubey MK. (2012, b). Biological effects of essential oils against rice weevil *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *J.Essen.l Oil bear. Pl.*, 15, 809-815.
- Chaudhary, S., Kanwar, R.K., Sehgal, A., Cahill, D.M., Barrow, C.J., Sehgal, R & Kanwar, J.R. (2017) Progress on *Azadirachta indica* based biopesticides in replacing synthetic toxic pesticides. *Front. Plant Sci.*, 8:610.
- Chogo, J.B., & Crank, G. (1981). Chemical composition and biological activity of the Tanzanian plant *Ocimum suave*. *Journal of Natural Products*, 44, 308-311.
- Choi, W., Lee, S.G., Park, H.M., & Ahn, Y.J. (2004). Toxicity of plant essential oils to *Tetranychus urticae* (Acari: Tetranychidae) and *Phytoseiulus persimilis* (Acari: Phytoseiidae). *Journal of Economic Entomology*, 97, 553-558.
- Chu, S.S., Liu, S.L., Jiang, G.H., & Liu, Z.L. (2010). Composition and toxicity of essential oil of *Illicium simonsii* Maxim (Illiciaceae) fruit against the maize weevils. *Records of Natural Products*, 4, 205-210.
- Cimanga, K., Kambu, K., Tona, L., Apers, S., De Bruyne, T., Hermans, N., Totte, J., Pieters, L., & Vlietinck, A.J. (2002). Correlation between chemical composition and antibacterial activity of essential oils of some aromatic medicinal plants growing in the democratic republic of Congo. *Journal of Ethnopharmacology*, no. 79, 213–220.
- Correa, Y. D. C. G., Faroni, L. R., Haddi, K., Oliveira, E. E. & Pereira, E. J. G. (2015). Locomotory and physiological responses induced by clove and cinnamon essential oils in the maize weevil *Sitophilus zeamais*. *Pestic. Biochem. Physiol.*, 125, 31–37.
- Dave, H., & Lediwane, L., (2012). A review on anthraquinones isolated from *Cassia* species and their applications. *Indian J. Nat. Prod. Resour.*, 3, 291–319.
- de Araújo, A. M. N. , Faroni, D.L.R., de Oliveira, J.V., Navarro, D.M.F., Barbosa, D.R.S., Breda, M.O., de França, S.M. (2017). Lethal and sublethal responses of *Sitophilus zeamais* populations to essential oils. *J. Pest Sci.*, 90, 2:589-600.

- Dehghani-Samani, A., Madreseh-Ghahfarokhi, S., Dehghani-Samani, A., & Pirali-Kheirabadi, K. (2015). Acaricidal and repellent activities of essential oil of *Eucalyptus globulus* against *Dermanyssus gallinae* (Acari: Mesostigmata). *J. Herb Med. Pharmacol.*, 4,3: 81-84.
- Diwan, R.K., & Saxena, R.C. (2010). Insecticidal property of flavinoid isolated from *Tephrosia purpuria*. *Int. J. Chem. Sci.*, 8, 2: 777-782
- Don-Perdo, K.M. (1996). Investigation of single and joint fumigant insecticidal action of citrus peel oil components. *J. Pest Sci.*, 46, 79-84.
- Elango, G., Rahuman, A.A., Bagavan, A., Kamaraj, C., Zahir, A.A., & Venkatesan, C. (2009). Laboratory study on larvicidal activity of indigenous plant extracts against *Anopheles subpictus* and *Culex tritaeniorhynchus*. *Parasitology research*, 104(6):1381-1388.
- El-Nahal, A.K.M., Schmidt, G.H., & Risha, E.M. (1989). Vapours of *Acorus calamus* oil – a space treatment for stored product insects. *Journal of Stored Products Research*, 25:211-216.
- El-Zemity, S., Hussien, R., Saher, F., & Ahmed, Z. (2006). Acaricidal activities of some essential oils and their monoterpenoidal constituents against house dust mite, *Dermatophagoides pteronyssinus* (Acari: Pyroglyphidae), *Journal of Zhejiang University Science B*, 7, 957–962.
- Elzen, G.W., & Hardee, D.D. (2003). United state department of agricultural-agricultural research on managing insect resistance to insecticides. *Pest Manag. Sci.*, 59, 770-776.
- Emam, A. M., Swelam, E.S., & Megally, N. Y. (2009). Furocoumarin and quinolone alkaloid with larvicidal and antifeedant activities isolated from *Ruta chalepensis* leaves. *Journal of Natural Products*, 2:10-22.
- Enan, E.E. (2005). Molecular and pharmacological analysis of an octopamine receptor from american cockroach and fruit fly in response to plant essential oils. *Archives of Insect Biochemistry and Physiology*, 59:161-171.

- Fradin, M.S., & Day, J.F. (2002). Comparative efficacy of insect repellents against mosquito bites. *The New England Journal of Medicine*, 347, 13–18.
- Gardulf, A., Wohlfart, I., & Gustafson, R. (2004). A prospective cross-over field trial shows protection of lemon *Eucalyptus* extract against tick bites. *J. Med. Entomol.*, 41, 1064-1067.
- Germinara, G. S., Distefano, M. G., Acutis, L. D., Pati, S., Delfne, S., Cristofaro, A. D., Rotundo, G. (2017). Bioactivities of *Lavandula angustifolia* essential oil against the stored grain pest *Sitophilus granaries*. *Bulletin of Insectology*, 70, 1: 129-138.
- Ghavami, M. B., Poorastgoo, F., Taghiloo, B., & Mohammadi, J. (2017). Repellency effect of essential oils of some native plants and synthetic repellents against human flea, *Pulex irritans* (Siphonaptera: Pulicidae). *J. Arthropod-Borne Dis.*, 11,1: 105–115.
- Ghoneim, K., & Hamadah, K. (2017). Antifeedant activity and detrimental effect of Nimbecidine (0.03% Azadirachtin) on the nutritional performance of Egyptian cotton leafworm *Spodoptera littoralis* Bois. (Noctuidae: Lepidoptera). *Bio Bulletin*, 31: 39-55,
- Giner, M., Avilla, J., Balcells, M., Caccia, S., & Smaghe, G. (2012). Toxicity of allyl esters in insect cell lines and in *Spodoptera littoralis* larvae. *Arch. Insect Biochem. Physiol.*, 79, 1:18-30.
- Goławska, S., & Łukasik, I. (2012). Antifeedant activity of luteolin and genistein against the pea aphid, *Acyrtosiphon pisum*. *J. Pest Sci.*, 85:443–450.
- Golawska, S., Kapusta, I., Łukasik I., & Wojcicka A. (2008). Effect of phenolics on the pea aphid, *Acyrtosiphon pisum* (Harris) population on *Pisum sativum* L. (Fabaceae). *Pestycydy/Pesticides*, (3-4), 71-77.
- Goławska, S., Łukasik, I., Goławski, A., Kapusta, I., & Janda, B. (2010) Alfalfa (*Medicago sativa* L.) apigenin glycosides and their effect on the pea aphid (*Acyrtosiphon pisum*). *Pol. J. Environ. Stud.*, 19:913–920.

- Goławska, S., Łukasik, I., Kapusta, I., & Janda, B. (2012). Do the contents of luteolin, tricetin, and chrysoeriol glycosides in alfalfa (*Medicago sativa* L.) affect the behavior of pea aphid (*Acyrtosiphon pisum*)? *Pol. J. Environ. Stud.*, 21:1613–1619.
- Goławska, S., Sprawka, I., Łukasik, I., & Goławski, A. (2014). Are naringenin and quercetin useful chemicals in pest-management strategies? *J. Pest Sci.*, 87:173–180
- Gould, K.S.; & Lister, C. (2006). *Flavonoid functions in plants. In Flavonoids: Chemistry, Biochemistry, and Applications*; CRC Press LLC: Boca Raton, FL, USA, pp. 397–443.
- Guedes, R. N. C., Oliveira, E. E., Guedes, N. M. P., Ribeiro, B. & Serrão, J. E. (2006). Cost and mitigation of insecticide resistance in the maize weevil, *Sitophilus zeamais*. *Physiol. Entomol.*, **31**, 30–38.
- Halliwell, B. & Gutteridge, J. M. C. (1999). *Free radicals in biology and medicine*, 3rd edn. Oxford University Press, UK.
- Hieu, T. T., Choi, W. S., Kim, S.I., Wang, M., & Ahn, Y. J. (2015). Enhanced repellency of binary mixtures of *Calophyllum inophyllum* nut oil fatty acids or their esters and three terpenoids to *Stomoxys calcitrans*. *Pest Manag. Sci.*, 71, 1213–1218.
- Ho S.H., Ma, Y., Goh, P.M., & Sim, K.Y. (1995). Star anise, *Illicium verum* Hook F., as a potential grain protectant against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* (Motsch.). *Post Harvest Biology and Technology*, 6, 341-347.
- Ho, S. H., Koh, L., Ma, Y., Huang, Y. & Sim, K. Y. (1996). The oil of garlic, *Allium sativum* L. (Amaryllidaceae), as a potential grain protectant against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Postharvest Biol. Technol.*, 9, 41–48.
- Hollingworth, R., Ahammadsahib, K., Gadelhak, G., & McLaughlin, J. (1994). New inhibitors of complex I of the mitochondrial electron transport chain with activity as pesticides. *Biochemical Society Transactions*, 22, 1:230-233.
- Hollingworth, R.M., Johnstone, E.M., & Wright, N. (1984). In: pesticide synthesis through rational approaches, Magee, PS, Kohn GK, Menn JJ (Eds), ACS symposium series No. 255: american chemical society, Washington, DC. 103-125.

- Houghton, P.J., Ren, Y., & Howes, M.J. (2006). Acetylcholinesterase inhibitors from plants and fungi. *Natural Product Reports*, 23, 2:181-199.
- Huang, Y., Lam, S.L., & Ho, S.H. (2000). Bioactivities of essential oils from *Elletaria cardamomum* (L.) Maton. to *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst). *Journal of Stored Products Research*, 36, 107-117.
- ICIPE. *International Centre of Insect Physiology and Ecology* (1997). *Vision and Strategic framework towards 2020*. ICIPE Science Press. Nairobi, Kenya.
- Ileke, K.D., & Olotuah, O.F. (2012). Bioactivity of *Anacardium occidentals* and *Allium sativum* powders and oils extracts against cowpea bruchid, *Callosobruchus maculatus* (Fab) (Coleoptera: Bruchidae). *International Journal of Biological Science*, 4, 1:96-103.
- Imms, A.D. (1964). *Outlines of Entomology*. 5th ed. pp. 224. Methuen. London, UK.
- Isman, M.B. (2006). Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annu. Rev. Entomol.*, 51, 45-66.
- Isman, M.B., & Machial, C.M. (2006). Pesticides based on plant essential oils: from traditional practice to commercialization. In: Rai, M., Carpinella, M.C. (Eds.), naturally occurring bioactive compounds. *Advances in Phytomedicine*, 3, 29-44.
- Jayakumar, M., Arivoli, S., Raveen, R., & Tennyson, S. (2017). Repellent activity and fumigant toxicity of a few plant oils against the adult rice weevil *Sitophilus oryzae* Linnaeus 1763 (Coleoptera: Curculionidae). *Journal of Entomology and Zoology Studies*, 5, 2: 324-335
- Kanat, M., Hakk, M., & Alma, M. (2003). Insecticidal effects of essential oils from various plants against larvae of pine processionary moth (*Thaum etopoeapityocampa* Schiff) (Lepidoptera: Thaumetopoeidae). *Pest Manag. Sci.*, 60, 173-177.
- Karunamoorthi, K., Girmay, A., & Hayleeyesus, S.F. (2014). Mosquito repellent activity of essential oil of Ethiopian ethnomedicinal plant against Afro-tropical malarial vector *Anopheles arabiensis*. *Journal of King Saud University of Science*, 26, 305-310.

- Kaufmann, C., & Briegel, H. (2004). Flight performance of the malaria vectors *Anopheles gambiae* and *Anopheles atroparous*. *Journal of Vector Ecology*, 29, 1:140- 153.
- Kim, S.I., Roh, J.Y., Kim, D.H., Lee, H.S., Ahn, Y.J. (2003). Insecticidal activities of aromatic plant extracts and essential oils against *Sitophilus oryzae* and *Callosobruchus chinensis*. *Journal of Stored Products Research*, 39, 293-303.
- Kimutai, A., Ngeiywa, M., Mula, M., Njagi, P.G.N., Ingonga, J., Nyamwamu, L.B., Ombati, C., & Ngumbi, P. (2017). Repellent effects of the essential oils of *Cymbopogon citratus* and *Tagetes minuta* on the sandfly, *Phlebotomus duboscqi*. *BMC Res Notes*, 10, 98.
- Carroll, J.F., Demirci, P., Kramer, M., Bernier, U.R., Agramonte, N.M., Baser, K.H.C., & Tabanca, N. (2017). Repellency of the *Origanum onites* L. essential oil and constituents to the lone star tick and yellow fever mosquito. *Natural Product Research*, 31, 18: 2192-2197.
- Jose, S., & Sujatha, K. (2017). Antifeedant activity of different solvent extracts of *Gliricidia sepium* against third instar larvae of *Helicoverpa armigera* (Hubner) (Lepidoptera : Noctuidae). *Int. J. Adv. Res. Biol. Sci.*, 4, 4: 201-204
- Kordali, S., Cakir, A., Mavi, A., Kilic, H., & Yildirim, A. (2005). Screening of chemical composition and antifungal and antioxidant activities of the essential oils from three Turkish artemisia species. *J. Agric. Food Chem.*, 53, 1408–1416.
- Koshier, E.L., & Sedy, K.A. (2001). *Effect of plant volatiles on the feeding and oviposition of Thripstabaci*. In: Marullo, Kound, L. Eds. *Thrips and Tospoviruses*, CSIRO, Australia, pp. 185-187, 2001.
- Koul, O., Waliai, S., & Dhaliwal, G.S. (2008). Essential oils as green pesticides: potential and constraints. *Biopesticides International*, 4, 1:63-84.
- Kraiss, H., & Cullen, E.M. (2008). Insect growth regulator effects of azadirachtin and neem oil on survivorship, development and fecundity of *Aphis glycines* (Homoptera: Aphididae) and its predator, *Harmonia axyridis* (Coleoptera: Coccinellidae). *Pest manag.sci.*, 64, 6:660-668.

- Kubo, I., & Kim, M. (1989). New insect growth inhibitory flavan glycosides from *Viscum tuberculatum* Tetrahedron. *Letters*, 28, 9: 921-924.
- Kumar, P., Bhadauria, T., & Mishra, J. (2015). Impact of application of insecticide quercetin/azadirachtin and chlorpyrifos on earthworm activities in experimental soils in Uttar Pradesh India. *Science Postprint*, 1, 2: e00044.
- Lee, B.H., Choi, W.S., Lee, S.E., & Park, B.S. (2001 a). Fumigant toxicity of essential oils and their constituent compounds towards the rice weevil, *Sitophilus oryzae* (L.). *Crop Protection*, 20, 317-320.
- Lee, B.H., Lee, S.E., Annis, P.C., Pratt, S.J., Park, B.S., & Tumaalii, F. (2002). Fumigant toxicity of essential oils and Monoterpenes against the red flour beetle, *Tribolium castaneum* Herbst. *Journal of Asia-Pacific Entomology*, 5, 2:237-240.
- Lee, S.E. (2000). Mosquito larvicidal activity of piperonaline, a piperidine alkaloid derived from long pepper, *Piper longum*. *J. Am. Mosq. Control Assoc.*, 16, 3:245-7.
- Lee, S.E., Lee, B.H., Choi, W.S., Park, B.S., Kim, J.G., & Campbell, B.C. (2001b). Fumigant toxicity of volatile natural products from Korean spices and medicinal plants towards the rice weevil, *Sitophilus oryzae* (L.). *Pest Manag. Sci.*, 57, 548-553.
- Li, H., Madden J.L., & Potts, B.M. (1996). Variation in volatile leaf oils of the Tasmanian *Eucalyptus* species II. Subgenus Symphyomyrtus. *Biochemical Systematics and Ecology*, 24, 547-569.
- Liao, M., Xiao, J. J., Zhou, L. J., Yao, X., Tang, F., Hua, R. M., Wu, X. W. & Cao, H. Q. (2017). Chemical composition, insecticidal and biochemical effects of *Melaleuca alternifolia* essential oil on the *Helicoverpa armigera*. *J. Appl. Entomol.*, 2017, 1-8.
- Liu, X., Chen, Q., Wang, Z., Xie, L., & Xu, Z. (2008). Allelopathic effects of essential oil from *Eucalyptus grandis*, *E. urophylla* on pathogenic fungi and pest insects. *Frontiers of Forestry in China*, 3, 232-236.

- Liu, Z.H., & Ho, S.H. (1999). Bioactivity of essential oil extracted from *Evodia rutracurpa* (Hook F. et Thomas) against the grain storage insects, *Sitophilus zeamais* (Motsch.) and *Tribolium castaneum* (Herbst.). *Journal of Stored Products Research*, 35, 317-328.
- Loko, L. Y., Alagbe, O., Dannon, E.A., Datinon, B., Orobiyi, A., Thomas-Odjo, A., Dansi, A., & Tamò, M. (2017). Repellent effect and insecticidal activities of *Bridelia ferruginea*, *Blighia sapida*, and *Khaya senegalensis* leaves powders and extracts against *Dinoderus porcellus* in infested dried yam chips. *Psyche*, 2017 Article ID 5468202, 18 pages.
- López, S.B., López, M.L., Aragón, L.M., Tereschuk, M.L., Slanis, A.C., Feresin, G.E., Zygodlo, J.A., & Tapia, A.A. (2011). Composition and anti-insect activity of essential oils from *Tagetes* Species (Asteraceae, Helenieae) on *Ceratitis capitata* Wiedemann and *Triatoma infestans* Klug. *Journal of Agricultural and Food Chemistry*, 59, 10: 5286–5292.
- Lu, F.C. (1995). A review of the acceptable daily intakes of pesticides assessed by the world health organization. *Regulatory Toxicology and Pharmacology*, 21, 351-364.
- Lucia, A., Audino, P.G., Seccacini, E., Licastro, S., Zerba, E., & Masuh, H. (2007). Larvicidal effect of *Eucalyptus grandis* essential oil and turpentine and their major components on *Aedes aegypti* larvae. *J. Am. Mosq. Control Assoc.*, 23, 299-303.
- Lucia, A., Toloza, A. C., Guzmán, E., Ortega, F., & Rubio, R. G. (2017), Novel polymeric micelles for insect pest control: encapsulation of essential oil monoterpenes inside a triblock copolymer shell for head lice control. *Peer J.*, 5:e3171
- Malek, M., & Parveen, B. (1989). Effect of insects infestation on the weight loss and viability of stored BE paddy. *Bangladesh Journal of Zoology*, 17, 1: 83-85.
- Mallikarjuna, N., Kranthi, K.R., Jadhav, D.R., Kranthi, S., & Chandra, S. (2004). Influence of foliar chemical compounds on the development of *Spodoptera litura* in interspecific derivatives of groundnut. *J. Appl. Entomol.*, 128, 321–328.
- Mann, R.S., & Kaufman, P.E. (2012). Natural product pesticides: their development, delivery and use against insect vectors. *Mini-reviews in Organic Chemistry*, 9, 185-202.

- Mathur, Y.K., Shankar, K., & Ram, S. (1985). Evaluation of some grain protectants against *Callosobruchus chinensis* (L.) on black gram. *Bulletin of Grain Technology*, 23, 253-259.
- Matsumura, F. (1985). Toxicology of insecticides (2ed.), Plenum Press, New York. PP., 11-43.
- Mikhael, A.A. (2011). Potential of some volatile oils in protecting packages of irradiated wheat flour against *Ephestia kuheniella* and *Tribolium castaneum*. *Journal of Stored Products Research*, 47, 4:357- 364.
- Min, L., Jin-Jing, X., Li-Jun, Z., Liu, Y., Xiang-Wei, W., Ri-Mao, H., Gui-Rong, W., & Hai-Qun, C. (2016). Insecticidal activity of *Melaleuca alternifolia* essential oil and RNA-seq analysis of *Sitophilus zeamais* transcriptome in response to oil fumigation. *Plos One*, 11, 12: e0167748.
- Moretti, M.D.L., Sanna-Passino, G., Demontis, S., & Bazzoni, E. (2000). Essential oil formulations useful as a new tool for insect pest control. *AAPS Pharm Sci. Tech.*, 3, 2:13.
- Morimoto, M., Kumeda, S., & Komai, K. (2000). Insect antifeedant flavonoids from *Gnaphalium affine*. *J. Agric. Food Chem.*, 48:1888–1891.
- Morrison, N.I., Franz, G., Koukidou, M., Miller, T.A., Saccone, G., Alpey, L.S., Beech, C.J., Nagaraju, J., Simmons, G.S., & Polito, L.C. (2010). Genetic improvements to the sterile insect technique for agricultural pests. *As. Pac. J. Mol. Biol. Biotech.*, 18, 2:275-295.
- Mullens, B.A., Reifenrath, W.G., & Butler, S.M. (2009). Laboratory trials of fatty acids as repellents or antifeedants against houseflies, horn flies and stable flies (Diptera: Muscidae). *Pest Manag. Sci.*, 65(12):1360-6.
- Navarro-Llopis, V., Vacas, S., Sanchis, J., Primo, J., & Alfaro, C. (2011). Chemosterilant bait stations coupled with sterile insect technique: An integrated strategy to control the mediterranean fruit fly (Diptera: Tephritidae). *Journal of economic entomology*, 104, 5:1647-1655.

- Negahban, M., Moharramipour, S., & Sefidkon, F. (2007). Fumigant toxicity of essential oil from *Artemisia sieberi* Besser against three stored product insects. *Journal of Stored Products Research*, 43, 2:123-128.
- Nerio, L.S., Olivero-Verbel, J., & Stashenko, E. (2009). Repellency activity of essential oils from seven aromatic plants grown in Colombia against *Sitophilus zeamais* Motschulsky (Coleoptera). *Journal of Stored Products Research*, 45, 212-214.
- Obeng-Ofori, D., & Amitaye, S. (2005). Efficacy of mixing vegetable oils with pirimiphos-methyl against the maize weevil, *Sitophilus zeamias* Motchulsky, in stored maize. *Journal of Stored Products Research*, 41, 1:57-66.
- Padin, S.B., Fuse, C., Urrutia, M.I., & DalBello, G.M. (2013). Toxicity and repellency of nine medicinal plants against *Tribolium castaneum* in stored wheat. *Bulletin of Insectology*, 66, 1:45-49.
- Papachristos, D.P., Karamanoli, K., Stamopoulos, D.C., & Menkissoglu-Spiroudi, U. (2004). The relationship between the chemical composition of three essential oils and their insecticidal activity against *Acanthoscelides obtectus* (Say). *Pest Manag. Sci.*, 60, 514–520.
- Papanastasiou, S.A., Bali, E.M.D., Ioannou, C.S., Papachristos, D.P., Zarpas, K.D., & Papadopoulos, N.T. (2017). Toxic and hormetic-like effects of three components of citrus essential oils on adult Mediterranean fruit flies (*Ceratitis capitata*). *PLoS ONE*, 12, 5: e0177837.
- Park, D.S., & Coats, J.R. (2002). Cyanogenic glycosides: Alternative insecticides? *The Korean Journal of Pesticide Science*, 6, 2, 51-57.
- Park, J., Y.Jeon, C.Lee, N. Chung & H.Lee (2017). Insecticidal toxicities of carvacrol and thymol derived from *Thymus vulgaris* Lin. against *Pochazia shantungensis* Chou & Lu., newly recorded pest. *Sci. Rep.*, 2017; 7: 40902.

- Plata-Rueda, A., Martínez, L. C., Santos, M. H. D., Fernandes, F. L., Wilcken, C. F., Soares, M. A., Serrão, J. E. & Zanuncio, J. C. (2017). Insecticidal activity of garlic essential oil and their constituents against the mealworm beetle, *Tenebrio molitor* Linnaeus (Coleoptera: Tenebrionidae). *Scientific Reports*, 7:46406.
- Pereira, S.G., Sanaveerappanavar, V.T., & Murthy, M.S. (2006). Geographical variation in the susceptibility of the diamond back moth *Plutella xylostella* L. to *Bacillus thuringiensis* products and acylurea compounds. *Pest Management*, 15, 26-26.
- Perumalsamy, H., Jang, M. J., Kim, J.R., Kadarkarai, M., & Ahn, Y.J. (2015). Larvicidal activity and possible mode of action of four flavonoids and two fatty acids identified in *Millettia pinnata* seed toward three mosquito species. *Parasites & Vectors*, 8, 237–244.
- Priestley, C.M., Burgess, I.F., & Williamson, E.M. (2006). Lethality of essential oil constituents towards the human louse, *Pediculus humanus* and its eggs. *Fitoterapia*, 77, 303-309.
- Prowse, M. G., Galloway, T. S. & Foggo, A. (2006). Insecticidal activity of garlic juice in two dipteran pests. *Agr. Forest Entomol.*, 8, 1–6.
- Pugazhvendan, S.R., Ross, P.R., & Elumalai, K. (2012). Insecticidal and repellent activities of four indigenous medicinal plants against stored grain pest, *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Asian Pacific Journal of Tropical Disease*, S16-S20.
- Pujiarti, R., & Fentiyanti, P.K. (2017). Chemical compositions and repellent activity of *Eucalyptus tereticornis* and *Eucalyptus deglupta* essential oils against *Culex quinquefasciatus* mosquito. *Thai Journal of Pharmaceutical Sciences*, 41, 1: 19–24.
- Qari, S. H., Nilly A.H., Abdel-Fattah, A.H., & Shehawy, A.A. (2017). Assessment of DNA damage and biochemical responses in *Rhyzopertha dominica* exposed to some plant volatile oils. *Journal of Pharmacology and Toxicology*, 12, 87-96
- Rajashekar, Y., Bakthavatsalam, N., Shivanandappa, T. (2012). Botanicals as grain protectants. *Psyche*, 2012, 1-13.

- Rahdari, T., & M. Hamzei (2017). Repellency Effect of essential oils of *Mentha piperita*, *Rosmarinus officinalis* and *Coriandrum sativum* on *Tribolium confusum* duval (Coleoptera: Tenebrionidae). *Chemistry Research Journal*, 2, 2:107-112.
- Rattan, R. S. (2010). Mechanism of action of insecticidal secondary metabolites of plant origin. *Crop Prot.*, 29, 913–920.
- Regnault-Roger, C., & Philogène, B.J.R. (2008). Past and current prospects for the use of botanicals and plant allelochemicals in integrated pest management. *Pharmaceutical Biology*, 46, 41–52.
- Regnault-Roger, C. (1997). The potential of botanical essential oils for insect pest control. *Integrated Pest Management Reviews*, 2, 25-34.
- Regnault-Roger, C., & Hamraoui, A. (1993). Influence d'huiles essentielles sur *Acanthoscelides obtectus* Say, bruche du haricot. *Acta Botanica Gallica*, 140, 217-222.
- Regnault-Roger, C., Vincent, C., & Arnason, J.T. (2012). Essential oils in insect control: low-risk products in a high-stakes world. *Annu. Rev. Entomol.*, 57:405-24.
- Rotimi, O.A., Chris, O.A., Olusola, O.O., Joshua, R., & Josiah, A.O. (2011). Bioefficacy of extracts of some indigenous Nigerian plants on the developmental stages of mosquito (*Anopheles gambiae*). *Jordan Journal of Biological Sciences*, 4, 4:237-242.
- Samuel, M., Oliver, S. V., Wood, O. R., M. Coetzee & Brooke, B. D. (2015). Evaluation of the toxicity and repellence of an organic fatty acids mixture (C8910) against insecticide susceptible and resistant strains of the major malaria vector *Anopheles funestus* Giles (Diptera: Culicidae). *Parasites & Vectors*, 8, 321.
- Santos, J.P., Maia, J.D.G., & Cruz, I. (1990). Damage to germination of seed corn caused by maize weevil (*Sitophilus zeamais*) and Angoumois grain moth (*Sitotroga cerealella*). *Pesquisa Agropecuaria Brasileira*, 25, 12: 1687-1692.

- Santos, P. C., Santos, V. H. M., Mecina, G. F., Andrade, A. R., Fegueiredo, P. A., Moraes, V.M. O., Silva, L. P., & Silva, R. M. G. (2016). Insecticidal activity of *Tagetes* sp. on *Sitophilus zeamais* Mots. *International Journal of Environmental & Agriculture Research*, 2, 4: 31-38.
- Schmidt, S., Tomasi, C., Pasqualini, E., & Ioriatti, C. (2008). The biological efficacy of pear ester on the activity of *Granulosis virus* for codling moth. *J. Pest Sci.*, 81: 29.
- Schoonhoven, A.V. (1978). The use of vegetable oils to protect stored beans from bruchid attack. *Journal of Economic Entomology*, 71, 2:254-256.
- Seyoum, A., Killeen, G.F., Kabiru, E.W., Knols, B.G.I., & Hassanali, A. (2003). Field efficacy of thermally expelled or live potted repellent plants against African malaria vectors in western Kenya. *Tropical Medicine & International Health*, 8, 1005–101,
- Shaalán, E.A.S, Canyon, D., Younes, M.W.F, Abdel-Wahab, H., & Mansour, A.H. (2005). A review of botanical phytochemicals with mosquitocidal potential. *Environment international*, 31, 8:1149-1166.
- Sharififard, M., Safdari, F., Siahpoush, A., Hamid H., & Kassiri, H. (2016). Evaluation of some plant essential oils against the brown-banded cockroach, *Supella longipalpa* (Blattaria: Ectobiidae): A mechanical vector of human pathogens. *Journal of Arthropod-Borne Diseases*, 10, 4: 528–537.
- Shelton, A.M., Zhao J.Z., & Roush, R.T. (2002). Economic, ecological, food safety, and social consequences of the deployment of B-transgenic plants. *Annual Review of Entomology*, 47, 845-88.
- Showler, A. T. (2017). Botanically based repellent and insecticidal effects against horn flies and stable flies (Diptera: Muscidae). *Journal of Integrated Pest Management*, 8,1: 15; 1–11
- Shukla, P., Vidyasgar, P. S. P. V., Aldosari, S. A., & Abdel-Azim, M. (2012). Antifeedant activity of three essential oils against the red palm weevil, *Rhynchophorus ferrugineus*. *Bulletin of Insectology*, 65, 1: 71-76.

- Silva, V. C.B., Ribeiro Neto, J. A., Alves, S. N., & Li, L. A. R.S. (2015). Larvicidal activity of oils, fatty acids, and methyl esters from ripe and unripe fruit of *Solanum lycocarpum* (Solanaceae) against the vector *Culex quinquefasciatus* (Diptera: Culicidae). *Revista da Sociedade Brasileira de Medicina Tropical.*, 48, 5:610-613.
- Simmonds, M.S. (2003). Flavonoid-insect interactions: Recent advances in our knowledge. *Phytochemistry*, 64, 21–30.
- Simmonds, M.S., & Stevenson, P.C. (2001). Effects of isoflavonoids from cicer on larvae of *Heliocoverpa armigera*. *J. Chem. Ecol.*, 27, 965–977.
- Singh, D., Siddiqui, M.S., & Sharma, S. (1989). Reproduction retardant and fumigant properties in essential oils against rice weevil (Coleoptera: Curculionidae) in stored wheat. *Journal of Economic Entomology*, 82:727-733.
- Singh, V. K. & Singh, D. K. (1996). Enzyme inhibition by allicin, the molluscicidal agent of *Allium sativum* L. (garlic). *Phytotherapy Res.*, 10, 383–386.
- Sithisut, D., Fields, P.G., & Chandrapathya, A. (2011). Contact toxicity, feeding reduction and repellency of essential oils from three plants from the ginger family (Zingiberaceae) and their major components against *Sitophilus zeamais* and *Tribolium castaneum*. *The Journal of Stored Products*, 104, 1445-54.
- Talukder, F., Islam, M., Hossain, M., Rahman, M., & Alam, M. (2004). Toxicity effects of botanicals and synthetic insecticides on *Tribolium castaneum* (Herbst) and *Rhyzopertha dominica* (F.). *Bangladesh J. Environ. Sci.*, 10, 2:365-371.
- Talukder, F.A. (2006). Plant products as potential stored-product insect management agents-A mini review. *Emir J Agric Sci.*, 18, 1:17-32.
- Tholl, D. (2006). Terpene synthases and the regulation, diversity and biological roles of terpene metabolism. *Curr Opin Plant Biol.*, 9, 297–304.

- Toloza, A.C., Zygodlo, J., Cueto, G.M., Biurrun, F., Zerba, E., & Piccolo, M.S. (2006). Fumigant and repellent properties of essential oils and component compounds against permethrin-resistant *Pediculus humanus capitis* (Anoplura: Pediculidae) from Argentina. *Journal of Medical Entomology*, 43, 5: 889-895.
- Tripathi, A.K., Prajapati, V., Aggarwal, K.K., Kumar, S., Kukreja, A.K., Dwivedi, S., & Singh, A.K. (2000). Effects of volatile oil constituents of *Mentha* species against stored grain pests, *Callosobruchus maculatus* and *Tribolium castaneum*. *Journal of Medicinal and Aromatic Plant Sciences*, 22, 549–556, 2000.
- Tripathi, A.K., Prajapati, V., Khanuja, S.P.S., & Kumar, S. (2003). Effect of d-limonene on three Stored-product beetles. *Journal of Economic Entomology*, 96, 990-995.
- Trivedi, A., Nayak, N., & Kumar, J. (2017). Fumigant toxicity study of different essential oils against stored grain pest *Callosobruchus chinensis*. *Journal of Pharmacognosy and Phytochemistry*, 6, 4: 1708-1711.
- Veal, L. (1996). The potential effectiveness of essential oils as a treatment for headlice, *Pediculus humanus capitis*. *Complementary Therapies in Nursing and Midwifery*, 2, 97-101.
- Velu, K., Elumalai, D., Hemalatha, P., Babu, M., Janaki, A., & Kaleena, P. K. (2015). Phytochemical screening and larvicidal activity of peel extracts of *Arachis hypogaea* against chikungunya and malarial vectors. *International Journal of Mosquito Research*, 2, 1: 01-08.
- Wachira, S. W., Omar, S., Jacob, J. W., Wahome, M., Alborn, H. T., Spring, D. R., Masiga, D. K., & Torto B. (2014). Toxicity of six plant extracts and two pyridine alkaloids from *Ricinus communis* against the malaria vector *Anopheles gambiae*. *Parasites & Vectors*, 7, 312.
- Waliwitiya, R., Kennedy, C., & Lowenberger, C. (2008). Larvicidal and oviposition altering activity of monoterpenoids, trans-anethole and rosemary oil to the yellow fever mosquito *Aedes aegypti* (Diptera: Culicidae). *Pest Manag. Sci.*, 65, 3:241-248.

- Watanabe, K., Shono, Y., Kakimizu, A., Okada, A., Matsuo, N., Satoh, A., & Nishimura, H. (1993). New mosquito repellent from *Eucalyptus camaldulensis*. *J. Agric. Food Chem.*, 41, 2164-2166.
- Weaver, D.K., Dunkel, F.V., Ntezurubanza, L., Jackson, L.L., & Stock, D.T. (1991). The efficacy of linalool, a major component of freshly-milled *Ocimum canum* Sims (Lamiaceae), for protection against postharvest damage by certain stored product Coleoptera. *Journal of Stored Products Research*, 27, 213-220.
- Wilke, A.B.B., Nimmo, D.D., John, O., Kojin, B.B., Capurro, M.L., & Marrelli, M.T. (2009). Mini-review: genetic enhancements to the sterile insect technique to control mosquito populations. *As. Pac. J. Mol. Biol. Biotech.*, 17, 3:65-74.
- Wimmer, Z., Alexandra, J.F.D.M. Floro, Zarevúcka, M., Wimmerová, M., Sello, G., & Orsini, F. (2007). Insect pest control agents: novel chiral butanoate esters (juvenogens) *Bioorg Med. Chem.*, 15, 18: 6037–6042.
- Wu, Y., Guo, S., Huang, D., Wang, C., Wei, J., Li, Z., Sun, J., Bai, J., Tian, Z., Wang, P., & Du, S. (2017). Contact and repellent activities of zerumbone and its analogues from the essential oil of *Zingiber zerumbet* (L.) Smith against *Lasioderma serricorne*. *J.oleo sci.*, 66, 4: 399-405.
- Yousef, H., EL-Lakwah, S. F., & EL Sayed, Y. A. (2013). Insecticidal activity of linoleic acid against *Spodoptera littoralis* (BOISD.). *Egypt. J. Agric. Res.*, 91, 2: 573.
- Zagrobelny, M., Bak, S., Rasmussen, A.V., Jørgensen, B., Naumann, C.M., & Møller, B.L. (2004). Cyanogenic glucosides and plant-insect interactions. *Phytochemistry*, 65, 3: 293–306.
- Zhang, W., Zhang, Z., Chen, Z., Liang, J., Geng, Z., Guo, S., Du, S., and Deng, Z. (2017). Chemical composition of essential oils from six *Zanthoxylum* Species and their Repellent activities against two stored-product insects. *J. Chem.*, 2017, Article ID 1287362, 7 pages.

Zhao, M. P., Liu, Q. Z., Liu Q., & Liu Z. L. (2017). Identification of larvicidal constituents of the essential oil of *Echinops grijsii* roots against the three species of mosquitoes. *Molecules*, 22, 205.

Zhao, N. N., Zhang, H., Zhang, X.C., Luan, X.B., Zhou, C., Liu, Q.Z., Shi, W.P., & Liu, Z.L. (2013). Evaluation of acute toxicity of essential oil of garlic (*Allium sativum*) and its selected major constituent compounds against overwintering *Cacopsylla chinensis* (Hemiptera: Psyllidae). *J. Econ. Entomol.*, 106, 1349–1354.

Zhu, J. J., Brewer, G. J., Boxler, D. J., Friesen, K., & Taylor, D. B. (2015). Comparisons of antifeedancy and spatial repellency of three natural product repellents against horn flies, *Haematobia irritans* (Diptera: Muscidae). *Pest Manag. Sci.*, 71, 1553–1560.

ACCEPTED MANUSCRIPT

## **ABOUT THE AUTHORS**

**Wafaa Hikal**, PhD, parasitology. Her research interests focus on parasitology and published peer-reviewed articles in the areas of parasitology and books, and collaborate in water borne parasites projects. She is a member of various scientific societies and editorial board member of some international journals.

**.Rowida Baeshen**, Assistant professor at the University of Tabuk, Saudi Arabia. Her research areas include entomology, and this paper is one of the main concerns of volatile oils application and one of the researcher's top priorities. She published peer-reviewed articles and book in the areas of entomology.

**Hussein Said-Al Ahl**, Professor at the National research Centre, and collaborates in medicinal and aromatic plants projects. His research interests focus on medicinal and aromatic plants. He is a member of various scientific societies and editorial board member of some international journals. Award-winning of the National Research Center Scientific Encouragement for agricultural sciences and published peer-reviewed articles and books.

## **PUBLIC INTEREST STATEMENT**

Provision of food has always been a challenge facing mankind. A major cornerstone in this challenge is the competition from insects. Insect pests inflict damage to humans, farm animals and crops. They are responsible for destroying one fifth of the world's total crop production annually. Considerable losses of agricultural products add a serious burden to people's daily life. There is an urgent need to move toward natural products as insecticides because it reduces negative impacts to human health and the environment, and we recommend the botanical insecticides to reduce the insect population and increase food production. Therefore, this study is useful for farmers, producers and exporters of crops and recommends the use of botanical insecticides as an alternative to chemicals or synthetic insecticides to eliminate insects and maintain crops and meet the risks of food shortages as well as the preservation of the environment and human and animal health.

ACCEPTED MANUSCRIPT