

## Human peptidoglycan recognition protein 1 in innate immunity

Jie Yang

### Accepted Manuscript Version

This is the unedited version of the article as it appeared upon acceptance by the journal. A final edited version of the article in the journal format will be made available soon.

As a service to authors and researchers we publish this version of the accepted manuscript (AM) as soon as possible after acceptance. Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). Please note that during production and pre-press, errors may be discovered which could affect the content.

© 2018 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

**Publisher:** Cogent OA

**Journal:** *Cogent Biology*

**DOI:** <http://doi.org/10.1080/23312025.2018.1450917>

## Human peptidoglycan recognition protein 1 in innate immunity

Jie Yang

Jie Yang <sup>1,2</sup>

<sup>1</sup>Central Laboratory, Danyang People's Hospital of Jiangsu Province, 2 Xinmin West Road, Danyang, Jiangsu Province, China

<sup>2</sup>Danyang Hospital Affiliated to Nantong University, Jiangsu Province, China

Corresponding author: yctcyangjie@163.com

**Abstract** Peptidoglycan recognition proteins (PGRPs) are innate immunity proteins that are conserved from insects to mammals. Related research results show the functions of PGLYRP1 in the innate immunity of neutrophils is to conducive to the killing of intracellular and extracellular bacteria. It is significant to make a study of human PGLYRP1 because neutrophils are a more dominant mechanism in host defense. Bactericidal activity of PGLYRP-1, PGLYRP-3, PGLYRP-4, and PGLYRP-3:4 for both Gram positive and negative bacteria requires  $Zn^{2+}$ . In addition, for killing of Gram negative,  $Zn^{2+}$  cannot be replaced by other cations, but for killing of Gram positive,  $Zn^{2+}$  can be partially replaced by  $Ca^{2+}$ , which have been proved. Then, the effect of PGLYRP1 on several human diseases have been reviewed. Taken together, these results indicate human *PGLYRP1* encodes an innate immunity protein that breaks down the structure of microbial cell wall, plays a role both in antibacterial defenses and inflammation and diseases.

**Key words** Inflammation; PGLYRP1; Disease; Innate immunity

## 1 Introduction

The innate immune system is a host defense mechanism, conserved from insects to mammalian evolutionarily, that mediates recognition and control of invading microorganisms [1]. PGRPs are innate immune proteins, and in some cases hydrolyze the peptidoglycans (PGNs) of bacterial cell walls [2]. When first insect and mammalian PGRPs are cloned in 1998, it is noticed that they all contain an amidase homology domain [3]. In 2000, a family of thirteen PGRP genes in *Drosophila* and a year later a family of four PGRPs in humans are identified [4]. They are initially named PGRP-S, PGRP-L, and PGRP-I $\alpha$  and PGRP-I $\beta$  (for ‘short’, ‘long’, or ‘intermediate’ transcripts, respectively) [5]. Later, the names for human PGRPs are changed by the Human Genome Organization Gene Nomenclature Committee to PGLYRP1, PGLYRP2, PGLYRP3, and PGLYRP4, respectively, and this nomenclature has been adopted for all mammalian PGRPs [6].

One mammalian PGRP, PGLYRP-2 is secreted from liver into blood [7], and is also induced by bacteria in epithelial cells [8]. The three remaining mammalian PGRPs are bactericidal or bacteriostatic proteins [9]. PGLYRP-1 is expressed primarily in the granules of polymorphonuclear leukocytes (PMNs), and PGLYRP-3 and PGLYRP-4 are expressed in the skin, eyes, mouth, intestinal tract, saliva, and protect the host against infections.

PGLYRP1 is approximately 200 amino acids, has a signal peptide and one PGRP domain, and a molecular mass of about 18-20 kDa. PGLYRP1 is highly expressed in the bone marrow in PMNs and their precursors [10], and the protein is almost exclusively present in the tertiary (secretory) granules, from which it could be released by exocytosis during phagocytosis.

## 2 Function of human PGLYRP1

PGRPs influence host-pathogen interactions not only through their antibacterial or peptidoglycan hydrolytic properties [11], but also through their pro-inflammatory and anti-inflammatory properties that are independent of their hydrolytic and antibacterial activities [12]. They maybe play a key role both in antibacterial defenses and several inflammatory diseases [13].

### 2.1 Bactericidal and bacteriostatic activity

Related research results show the functions of human PGLYRP1 in the innate immunity of neutrophils is to conducive to the killing of intracellular and extracellular

bacteria [14]. Although in initial studies purified human PGLYRP-1 are only bacteriostatic, the latest results have demonstrated that human PGLYRP-1 is bactericidal and that the bactericidal activity requires  $\text{Ca}^{2+}$ ; the earlier preparations are not bactericidal because they did not contain  $\text{Ca}^{2+}$  [15,16].

Subsequently, some studies have demonstrated that human PGLYRP-1, PGLYRP-3, PGLYRP-4, and PGLYRP-3:4 have  $\text{Zn}^{2+}$ -dependent bactericidal activity against both Gram positive and Gram negative bacteria at physiologic  $\text{Zn}^{2+}$  concentrations found in serum, sweat, saliva, and other body fluids. The requirement for  $\text{Zn}^{2+}$  can only be partially replaced by  $\text{Ca}^{2+}$  for killing of Gram positive bacteria but not for killing of Gram negative bacteria [17]. This  $\text{Zn}^{2+}$  dependence explains why in their previous experiments PGLYRPs purified in the presence of  $\text{Ca}^{2+}$  (without  $\text{Zn}^{2+}$ ) are not bactericidal for Gram negative bacteria and were only bactericidal for some Gram positive bacteria [18].

## 2.2 Role in inflammation and disease

Related studies have suggested that the PGLYRP1 dimer in human serum and polymorphonuclear cells is detected, from where it is secreted after degranulation; these cells being a possible source of serum PGLYRP1; these cells being a possible source of serum PGLYRP1 [19]. In addition, all PGRP protect cells from PGN-induced apoptosis. PGRP increase THP-1 cell proliferation and enhance activation by PGN. PGLYRP1-PGN complexes increase the membrane expression of CD14, CD80 and CD86, and enhance secretion of interleukin-8, interleukin-12 and tumor necrosis factor- $\alpha$ , but reduce interleukin-10, clearly inducing an inflammatory profile.

### 2.2.1 Inflammatory bowel disease (IBD)

IBD is a chronic inflammatory condition of the gastrointestinal tract. Triggering receptor expressed on myeloid cells 1 (TREM-1) is expressed on neutrophils and most monocytes or macrophages constitutively and potently amplifies inflammation [20]. Recently, PGLYRP-1 is shown to be the ligand of TREM-1 [21]. PGLYRP-1 is an antimicrobial peptide stored in neutrophil granules with PGN-binding activity [22]. Neutrophil degranulation releases PGLYRP-1 that, multimerized with itself or complexed with PGN, potently activates TREM-1 causing pro-inflammatory cytokine release. As PGN is a cell wall component of all bacteria, activation of TREM-1 cells by PGLYRP-1 or PGN complexes in bacteria-rich environments potently amplifies inflammation.

Here, the ability of an anti-TREM-1 antibody to dampen the release of pro-

inflammatory cytokines by colon lamina propria cells (LPCs) from patients with IBD is investigated and correlated with PGLYRP-1 levels. Moreover, PGLYRP-1+ myeloperoxidase is a potential biomarker for predicting the effect of anti-TREM-1 therapy [23]. These studies open the possibility for a new treatment for IBD and offer insight into PGLYRP-1, combined with a neutrophil marker such as MPO, as biomarkers to predict patients who would benefit from anti-TREM-1 therapy.

### 2.2.2 *ST-elevation myocardial infarction (STEMI)*

Acute myocardial infarction (AMI) is largely categorized into two categories: ST-segment-elevation myocardial infarction (STEMI) and non-ST-segment-elevation myocardial infarction (NSTEMI) [24]. Among the most easily accessible bio-fluids is the whole blood, containing leukocytes with informative transcripts used in their first line of immune defense and sentinels for many disease processes [25]. Using peripheral blood in clinical applications can also provide early and accurate information before development of the disease [26]. Accordingly, it is the potential method to be informative in disease status and of the underlying diverse disease mechanisms by the blood gene expression profiling [27].

Blood gene expression profiling reflects the status of diseases, and characteristic molecular signature provides a novel window on gene expression preceding acute coronary events [28]. Fortunately, the correlation results have indicated that PGLYRP1, interleukin-1 receptor-associated kinase 3 (IRAK3) and viral nervous necrosis 3 (VNN3) are more specific and sensitive diagnostic biomarkers for STEMI than traditional creatine kinase-MB (CK-MB) or troponin by a simple ELISA method. Ironically, they are the most sensitive STEMI biomarkers, none has been reported for cardiovascular disease or cardiac markers.

### 2.2.3 *Atherosclerosis*

The thiazolidinedione medications (TZDs), pioglitazone and rosiglitazone, improve a number of inflammatory markers associated with cardiovascular disease (CVD) and have favorable effects on imaging intermediates of atherosclerosis including carotid intima-media thickness [29], vascular inflammation, restenosis following stent implantation [30], and de novo coronary artery disease progression. Although rosiglitazone favorably affects myriad intermediate markers of atherosclerosis [31], it appears to increase myocardial infarction (MI) risk.

Some studies have found that rosiglitazone has a unfavorable effects on three novel inflammatory biomarkers previously shown to independently associate with

atherosclerosis (LT $\beta$ R, PGLYRP-1, and CCL23) and a favorable effect on another novel biomarker, sRAGE, which previously studies have been shown to be inversely associated with atherosclerosis [32].

PGN is detectable in varying levels in the circulation and has been identified by immunohistochemical staining in human atherosclerotic specimens. PGN may promote localized inflammation in non mucosal sites by stimulating the production of PGRPs. Among four known human sub-types of PGRP, PGLYRP-1 is expressed primarily in PMN granules, likely functioning as an antibacterial protein. Plasma levels of PGLYRP-1 may represent the systemic response to bacterial exposure, possibly underpinning the observed associations between bacterial infection and exposure with coronary heart disease.

#### 2.2.4 Rheumatoid arthritis (RA)

RA is a chronic inflammatory and autoimmune disease characterized by inflammation of the synovial membrane leading to the destruction of affected joints. The study of polymorphisms of genes differentially expressed may lead to the identification of putative causal genetic variants in multifactorial diseases such as RA. The list of genes to explore is established on the basis of the differential expression in RA vs controls, and included four up-regulated genes (*S100A8*, *PGLYRP1*, *RNASE2*, and *LY96*) and two down-regulated genes (*RUNX3* and *IL2RB*) [33]. The hypothesis is that the differentially expressed genes are associated with RA, using family-based methodology.

Recent studies have shown that PGRPs influence host-pathogen interactions through their pro-inflammatory or anti-inflammatory properties, which are independent of their antibacterial activities [34], modulating the balance between inflammatory T-helper (Th)17 cells and regulatory T cells. Taking into consideration the potential role of bacteria in the pathogenesis of RA, an up-regulation of *PGLYRP1* in the context of the *HLA-DRB1* shared epitope (SE) may influence the onset of the disease affecting inflammatory circuits, cytokine production, and possibly antigen presentation.

#### 2.2.5 Skin melanoma and renal carcinoma

During the last decade novel approaches for cancer treatment have been developed. Anti-tumor vaccination is considered to be one of the most promising of these. One of the possible ways to induce specific immune response is the use of tumor rejection antigens discovered during recent years [35,36]. Clinically important results of

vaccinotherapy are achieved in patients with melanoma and renal carcinoma in a number of studies. The results with this treatment are comparable to chemotherapy and immunotherapy.

Clinically important results of vaccinotherapy are achieved in patients with melanoma and renal carcinoma in a number of studies. The results with this treatment are comparable to chemotherapy and immunotherapy [37]. Recently, a phase I/II trial has been undertaken to define the feasibility, safety and anti-tumor effects of the autologous vaccine prepared by transferring *PGLYRP1* gene into malignant melanoma and renal cell carcinoma cells [38]. Related findings have indicated that vaccinotherapy of patients with skin melanoma and renal carcinoma with the *PGLYRP1*-modified autologous tumor cells is safe and do not have any significant side-effects. Although there is no clear correlation between changes in clinical and immunological status, it is possible to define the tendency of the vaccine to enforce the antigen-specific immune response. Further studies are required to determine whether promising effects on immune activation will result in an actual clinical benefit for patients with malignant melanoma and renal cell carcinoma.

### 3 Conclusions

In conclusion, human PGLYRP1 plays a role in innate immunity in the context of neutrophils by contributing to the killing of intracellular and extracellular bacteria. PGLYRP-1 have  $Zn^{2+}$ -dependent bactericidal activity against both Gram positive and Gram negative bacteria. Besides, the requirement for  $Zn^{2+}$  can only be partially replaced by  $Ca^{2+}$  for killing of Gram positive bacteria but not for killing of Gram negative bacteria, and also influence host-pathogen interactions through their pro-inflammatory and anti-inflammatory properties.

Recent findings indicate that the adaptive arm of immunity is governed by the innate immunity mechanisms that control co-stimulatory signaling of antigen-presenting cells (APCs) [39]. The combination of cytokines effecting both innate and acquired immune response taken together with tumor antigens may significantly improve overall efficiency of anti-tumor vaccination.

Since the PGLYRP1 dimer could be detected in human serum, and PGLYRP1–PGN complexes enhance the inflammatory response, it suggests that PGLYRP1 dimer may be a potentially serological markers during detecting human inflammation, and some human diseases. Although some functions of human PGLYRP1 have been identified, it is still possible that human PGLYRP1 have other unidentified functions, because many mammalian proteins have evolved to have multiple functions. In addition, it is still not clear how PGLYRP1 affects the cytokines in human various

diseases.

**Conflict of interest**

The author declares that there is no conflict of interest hampering the publication of this manuscript.

ACCEPTED MANUSCRIPT

## References

1. An JH, Kurokawa K, Jung DJ et al (2013) Human SAP is a novel peptidoglycan recognition protein that induces complement-independent phagocytosis of *Staphylococcus aureus*. *Journal of immunology* 191: 3319-3327
2. Nylund KM, Ruokonen H, Sorsa T et al (2017) Association of the Salivary Triggering Receptor Expressed on Myeloid Cells/ its Ligand Peptidoglycan Recognition Protein 1 Axis With Oral Inflammation in Kidney Disease. *Journal of periodontology*: 1-17
3. Kang D, Liu G, Lundstrom A et al (1998) A peptidoglycan recognition protein in innate immunity conserved from insects to humans. *Proceedings of the National Academy of Sciences of the United States of America* 95: 10078-10082
4. Chen K, Zhou L, Chen F et al (2016) Peptidoglycan recognition protein-S5 functions as a negative regulator of the antimicrobial peptide pathway in the silkworm, *Bombyx mori*. *Developmental and comparative immunology* 61: 126-135
5. Choe KM, Lee H, Anderson KV (2005) *Drosophila* peptidoglycan recognition protein LC (PGRP-LC) acts as a signal-transducing innate immune receptor. *Proceedings of the National Academy of Sciences of the United States of America* 102: 1122-1126
6. Cong M, Song L, Qiu L et al (2009) The expression of peptidoglycan recognition protein-S1 gene in the scallop *Chlamys farreri* was enhanced after a second challenge by *Listonella anguillarum*. *Journal of invertebrate pathology* 100: 120-122
7. Kashyap DR, Kuzma M, Kowalczyk DA et al (2017) Bactericidal peptidoglycan recognition protein induces oxidative stress in *Escherichia coli* through a block in respiratory chain and increase in central carbon catabolism. *Molecular microbiology* 105: 755-776
8. Kashyap DR, Wang M, Liu LH et al (2011) Peptidoglycan recognition proteins kill bacteria by activating protein-sensing two-component systems. *Nature medicine* 17: 676-683
9. Kibardin A, Karpova T, Sapenko T et al (2006) Mammalian peptidoglycan recognition protein TagL inhibits *Listeria monocytogenes* invasion into epithelial cells. *FEMS immunology and medical microbiology* 46: 284-290
10. Persson C, Oldenvi S, Steiner H (2007) Peptidoglycan recognition protein LF: a negative regulator of *Drosophila* immunity. *Insect biochemistry and molecular biology* 37: 1309-1316
11. Yang PJ, Zhan MY, Ye C et al (2017) Molecular cloning and characterization of a short peptidoglycan recognition protein from silkworm *Bombyx mori*. *Insect molecular biology* 26: 665-676
12. Wang N, Hirata A, Nokihara K et al (2016) Peptidoglycan microarray as a novel tool to explore protein-ligand recognition. *Biopolymers* 106: 422-429
13. Wang ZZ, Shi M, Huang YC et al (2016) A peptidoglycan recognition protein acts in whitefly (*Bemisia tabaci*) immunity and involves in Begomovirus acquisition. *Scientific reports* 6: 37806
14. Cho JH, Fraser IP, Fukase K et al (2005) Human peptidoglycan recognition protein S is an effector of neutrophil-mediated innate immunity. *Blood* 106: 2551-2558
15. Wang ZM, Li X, Cocklin RR et al (2003) Human peptidoglycan recognition protein-L is an N-acetylmuramoyl-L-alanine amidase. *The Journal of biological chemistry* 278: 49044-49052

16. Gelius E, Persson C, Karlsson J et al (2003) A mammalian peptidoglycan recognition protein with N-acetylmuramoyl-L-alanine amidase activity. *Biochemical and biophysical research communications* 306: 988-994
17. Wang M, Liu LH, Wang S et al (2007) Human peptidoglycan recognition proteins require zinc to kill both gram-positive and gram-negative bacteria and are synergistic with antibacterial peptides. *Journal of immunology* 178: 3116-3125
18. Lu X, Wang M, Qi J et al (2006) Peptidoglycan recognition proteins are a new class of human bactericidal proteins. *The Journal of biological chemistry* 281: 5895-5907
19. De Marzi MC, Todone M, Ganem MB et al (2015) Peptidoglycan recognition protein-peptidoglycan complexes increase monocyte/macrophage activation and enhance the inflammatory response. *Immunology* 145: 429-442
20. Bouchon A, Dietrich J, Colonna M (2000) Cutting edge: inflammatory responses can be triggered by TREM-1, a novel receptor expressed on neutrophils and monocytes. *Journal of immunology* 164: 4991-4995
21. Read CB, Kuijper JL, Hjorth SA et al (2015) Cutting Edge: identification of neutrophil PGLYRP1 as a ligand for TREM-1. *Journal of immunology* 194: 1417-1421
22. Stadnicki A, Colman RW (2003) Experimental models of inflammatory bowel disease. *Archivum immunologiae et therapiae experimentalis* 51: 149-155
23. Brynjolfsson SF, Magnusson MK, Kong PL et al (2016) An Antibody Against Triggering Receptor Expressed on Myeloid Cells 1 (TREM-1) Dampens Proinflammatory Cytokine Secretion by Lamina Propria Cells from Patients with IBD. *Inflammatory bowel diseases* 22: 1803-1811
24. Al Shehri MA, Youssef AA (2016) Acute myocardial infarction with multiple coronary thromboses in a young addict of amphetamines and benzodiazepines. *Journal of the Saudi Heart Association* 28: 180-184
25. Alkuraishy HM, Al-Gareeb AI (2015) New Insights into the Role of Metformin Effects on Serum Omentin-1 Levels in Acute Myocardial Infarction: Cross-Sectional Study. *Emergency medicine international* 2015: 283021
26. Andjic M, Spiroski D, Ilic Stojanovic O et al (2015) Effects of short-term exercise training in patients following Acute Myocardial Infarction treated with Primary Percutaneous Coronary Intervention. *European journal of physical and rehabilitation medicine*:
27. Bilal M, Haseeb A, Sher Khan MA (2015) Intracoronary infusion of Wharton's jelly-derived mesenchymal stem cells: a novel treatment in patients of acute myocardial infarction. *JPMA The Journal of the Pakistan Medical Association* 65: 1369
28. Andrechuk CR, Ceolim MF (2015) High risk for obstructive sleep apnea in patients with acute myocardial infarction. *Revista latino-americana de enfermagem* 23: 797-805
29. Milasan A, Ledoux J, Martel C (2015) Lymphatic network in atherosclerosis: the underestimated path. *Future science OA* 1: FSO61
30. Hong J, Maron DJ, Shirai T et al (2015) Accelerated atherosclerosis in patients with chronic inflammatory rheumatologic conditions. *International journal of clinical rheumatology* 10: 365-381
31. Ling Y, Jiang J, Gui M et al (2015) Thyroid Function, Prevalent Coronary Heart Disease, and Severity of Coronary Atherosclerosis in Patients Undergoing Coronary Angiography. *International journal of endocrinology* 2015: 708272

32. Lubrano V, Venturi E, Balzan S et al (2015) Impact Of Risk Factor for Atherosclerosis on Microvascular Endothelial Function: An In Vitro Study. *Theoretical biology forum* 108: 75-88
33. Fodil M, Teixeira VH, Chaudru V et al (2015) Relationship between SNPs and expression level for candidate genes in rheumatoid arthritis. *Scandinavian journal of rheumatology* 44: 2-7
34. Rohatgi A, Ayers CR, Khera A et al (2009) The association between peptidoglycan recognition protein-1 and coronary and peripheral atherosclerosis: Observations from the Dallas Heart Study. *Atherosclerosis* 203: 569-575
35. Sharapova TN, Ivanova OK, Soshnikova NV et al (2017) Innate Immunity Protein Tag7 Induces 3 Distinct Populations of Cytotoxic Cells That Use Different Mechanisms to Exhibit Their Antitumor Activity on Human Leukocyte Antigen-Deficient Cancer Cells. *Journal of innate immunity* 9: 598-608
36. Yashin DV, Ivanova OK, Soshnikova NV et al (2015) Tag7 (PGLYRP1) in Complex with Hsp70 Induces Alternative Cytotoxic Processes in Tumor Cells via TNFR1 Receptor. *The Journal of biological chemistry* 290: 21724-21731
37. Old LJ (1996) Immunotherapy for cancer. *Scientific American* 275: 136-143
38. Moiseyenko VM, Danilov AO, Baldueva IA et al (2005) Phase I/II trial of gene therapy with autologous tumor cells modified with tag7/PGRP-S gene in patients with disseminated solid tumors: miscellaneous tumors. *Annals of oncology : official journal of the European Society for Medical Oncology* 16: 162-168
39. Schnare M, Barton GM, Holt AC et al (2001) Toll-like receptors control activation of adaptive immune responses. *Nature immunology* 2: 947-950

Peptidoglycan recognition proteins (PGRPs) are innate immunity proteins that are conserved from insects to mammals. Related research results show the functions of PGLYRP1 in the innate immunity of neutrophils is to conducive to the killing of intracellular and extracellular bacteria. It is significant to make a study of human PGLYRP1 because neutrophils are a more dominant mechanism in host defense. This review highlights the role of PGLYRP1 in human immune responses. It focuses on an interesting an underappreciated family of pathogen recognition receptors, and attempts to show how PGLYRP1 could contribute to important human diseases, such as inflammatory bowel disease, ST-elevation myocardial infarction, atherosclerosis, rheumatoid arthritis, skin melanoma and renal carcinoma and so on.

Peptidoglycan recognition proteins (PGRPs) are innate immunity proteins that are conserved from insects to mammals. It is significant to make a study of human PGLYRP1 because neutrophils are a more dominant mechanism in host defense. Human PGLYRP1 plays a role in innate immunity in the context of neutrophils by contributing to the killing of intracellular and extracellular bacteria, and also influence host-pathogen interactions through their pro-inflammatory and anti-inflammatory properties. Since the PGLYRP1 dimer could be detected in human serum, and PGLYRP1-PGN complexes enhance the inflammatory response, it suggests that PGLYRP1 dimer may be a potentially serological markers during detecting human inflammation, and some human diseases.