



Received: 06 October 2016
Accepted: 17 November 2016
First Published: 29 November 2016

*Corresponding author: Julie Zuniga,
School of Nursing, The University of
Texas at Austin, 1700 Red River St.,
Austin, TX 78701, USA
E-mail: jzuniga@nursing.utexas.edu

Reviewing editor:
Albert Lee, The Chinese University of
Hong Kong, Hong Kong

Additional information is available at
the end of the article

PUBLIC HEALTH | RESEARCH ARTICLE

Effectiveness of using picture-based health education for people with low health literacy: An integrative review

Jungmin Park¹ and Julie Zuniga^{1*}

Abstract: *Background:* Level of health literacy affects health status. The purpose of this study was to examine and evaluate studies related to picture-based health education materials for people with low health literacy. *Methods:* Articles from 2001 through October 2015 were searched in the PubMed, CINAHL, and ERIC databases on the web, followed by manual searches of the references in those articles. *Results:* Of the 11 studies that met the inclusion criteria for this review, 10 showed a positive effect of increasing the health learning abilities of people with low health literacy using picture-based health education. Medication adherence was targeted most often. The majority of the studies measured health literacy using the Short Test of Functional Health Literacy in Adults, but the health literacy measurements in picture-based health education were nonetheless inconsistent. *Discussion:* Use of picture-based health education for people with low health literacy was limited. However, the majority of the studies did indicate a positive effect on learning.

Subjects: Education; Health and Social Care; Medicine, Dentistry, Nursing & Allied Health

Keywords: picture-based health education; low health literacy; cartoon; pictograph(s); illustration; integrative review

1. Introduction

Health literacy has been defined as “the degree to which individuals have the capacity to obtain, process, and understand basic health information and services needed to make appropriate health decisions” (Nielsen-Bohman, NetLibrary, & Institute of Medicine (US) Committee on Health

ABOUT THE AUTHORS

Jungmin Park is a doctoral candidate in the University of Texas at Austin, School of Nursing. Her research interest includes disease self-management by people in the HIV-infected population who have co-morbid conditions. Her dissertation focuses on retention in care for elderly people with HIV and diabetes. Jungmin Park currently serves as a research assistant on a study regarding self-management education of diabetes in HIV-infected people under the supervision of Dr Julie Zuniga, an assistant professor in the University of Texas at Austin, School of Nursing. Also, Jungmin is currently conducting a qualitative study investigating type two diabetes self-management education using pictography with low health literacy funded by the Sigma Theta Tau, Epsilon Theta Tau.

PUBLIC INTEREST STATEMENT

Low health literacy affects to negative health outcomes. The purpose of this manuscript is to integrate studies related to picture-based health education materials for people with low health literacy. In this paper, we used the online database and searched articles manually from 2001 to October 2015 under the PRISMA guideline. Ten articles indicate that picture-based health education materials helps to increasing the health learning abilities of people with low health literacy.

Literacy, 2004). In the US, 36% of adults have no more than basic health literacy (Kutner, Greenburg, Jin, & Paulsen, 2006). Most of those with low health literacy are elderly (over 65 years old), belong to minority groups, are refugees or immigrants, have a low educational level, have a low socioeconomic status, or are nonnative speakers (U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion, 2010).

People with low health literacy have many difficulties in managing their health problems (Arcia, Bales, & William Brown, 2013; Paasche-Orlow, Parker, Gazmararian, Nielsen-Bohlman, & Rudd, 2005) and as a result experience longer hospitalizations and more frequent use of emergency care. This may be due in part to difficulty accessing the correct medication, and understand labels and health messages (Arcia et al., 2013; Berkman, Sheridan, Donahue, Halpern, & Crotty, 2011). Those with limited health literacy are therefore more likely to become ill than those with higher health literacy (Garcia-Retamero, Okan, & Cokely, 2012).

Health education materials, however, are not always appropriately written for those with low health literacy, and the materials that do take low health literacy into account are limited. Health education materials are often written for a 10th-grade reading level (Davis et al., 1993; Doak, Doak, Friedell, & Meade, 1998) but the reading ability of people with low health literacy can be below the 5th-grade level (Kirsch & Educational Testing Service, & National Center for Education Statistics, 1993). One way to address this health knowledge gap is to use picture-based educational materials.

The Mayer's cognitive theory of multimedia learning (CTML) insisted three principles: (1) people process information through auditory and visual channels, (2) each channel has a limited amount of memory, and (3) learning consists of actively filtering and sorting information (Mayer & Ebooks Corporation, 2009). Given these principles, multimedia, which may be defined as "the combination of text and pictures," should help improve an individual's capacity to learn (Sorden, 2012), and the CTML is indeed based on the idea that learners can improve their learning abilities when they are presented with pictures as well as words (Choi, 2015; Mayer & Ebooks Corporation, 2009). Thus, the combination of pictures and words increases learning effectiveness (Mayer, 1947/2014; Sorden, 2012).


Following the Mayer's CTML theory, pictographs may increase the effectiveness of educational materials for those who have low literacy (Garcia-Retamero et al., 2012) especially for complex health information (Austin, Matlack, Dunn, Kesler, & Brown, 1995; Delp & Jones, 1996). However, pictographs encompass many different picture-based educational materials in a diverse health education fields, such as simple black and white line drawings, pictures, and simple drawings with simple text. The purpose of this integrative review is to synthesize studies associated with pictographic education designed for people with low health literacy, describe the characteristics of pictographs use, assess the tools used to measure health literacy in picture-based health education studies, and integrating the effectiveness of picture-based health education for people with low health literacy.

2. Methods

2.1. Search strategy

For this study, we systematically reviewed articles found with two main search methods: (1) electronic searches using key words in library databases, and (2) subsequent manual searches for related references. On the Internet, we searched three databases: PubMed, the Cumulative Index of Nursing and Allied Health Literature (CINAHL), and the Education Resources Information Center (ERIC). Search terms were as follows: *photo novella*, *education*, *cartoon*, *webtoon*, *pictograph*, *pictogram*, *illustration*, *picture*, *visualization*, *health*, *nonliterate*, and *literacy*. The research on picture-based materials for health education is limited, so we did not limit the time period for the review. Through an ancestry search, additional research studies were identified by searching reference lists that had inclusion criteria and purposes that were related to those of this review. The search strategy and data extraction are illustrated in Figure 1.

Figure 1. Search flow chart using <http://prisma.thetacollaborative.ca>.



2.2. Inclusion and exclusion criteria

This study was guided by inclusion and exclusion criteria: (1) English-based studies, (2) studies related to low health literacy, (3) studies use picture-based health education, (4) study participants are adults (over 18 years old), (5) studies are not reviews or meta-analyses, and (6) studies are not dental related.

2.3. Data extraction

Initially, the web-based search identified 70 articles in PubMed, CINAHL, and ERIC. Manual search of the references in those articles found an additional 22 articles, for a total of 92. After screening the 92 articles by title, key words, and abstract, 44 articles were excluded because they were editorials or presented only abstracts, were duplicates, were not written in English, applied to dental health of youth, were unrelated to health care, or did not consider participants' levels of health literacy. Full texts of the remaining 48 articles were then read. Review of those 48 articles led to exclusion of 20 articles from the web-based search because (1) the participants were not adults (over the age of 18), or (2) the articles were reviews or meta-analyses. A total of 17 articles from the manual search were also excluded after applying the same inclusion and exclusion criteria.

3. Results

3.1. Study samples and settings

The web-based search and the manual search yielded a final total of 11 articles (six from the web-based search, five from manual search) that met the inclusion criteria (see Table 1). Although there was no limit on the time period for the studies, all 11 articles were published from 2001 to 2015. The studies' sample sizes ranged from 6 to 500, and the samples included a broad range of participants. The final 11 articles included six cross-sectional descriptive studies (Choi, 2011, 2012, 2013; Chuang, Lin, Wang, & Cham, 2010; DeWalt et al., 2004; Kripalani et al., 2007) and seven interventions (Choi, 2015; DeWalt et al., 2004; Dowse & Ehlers, 2001, 2005; Kalichman et al., 2013; Kripalani et al., 2007; Mansoor & Dowse, 2003) with two of the articles presenting both descriptive studies and interventions (DeWalt et al., 2004; Kripalani et al., 2007) (see Table 2). The participants' ethnicities were diverse: Hispanic/Latino, White, African American, and Asian. The most common educational level

Table 1. Characteristics of picture-based health education strategies designed for patients with low health literacy

Study	Purpose	Setting	Sample characteristics	Design	Disease focus	Literacy measures	Key findings
Choi (2011)	To develop and validate pictograph-based discharge instructions for older adults after hip replacement surgery	United States	Experts assess appropriateness, accuracy, relevance N = 5 Age range: 39–51 years Clinical experience: 5–15 years 2 White, 2 AA, 1 Asian	Tool development and evaluation	Hip replacement surgery	Not appropriate	Pictographs (simple line drawings) assist engagement and are easy to understand for low-literate immigrant refugees in discharge care actions
Choi (2012)	To develop breast healthcare instructions using pictographs	United States	Immigrant women with low health literacy N = 6 Age range: 37–55 years Living in US: ≥10 years (83.3%) Edu. level: high school (66.7%) Hispanic/Latino (66.7%) Asian (16.7%)	Pilot study Tool development and evaluation Cross-sectional study Interviews	Breast health care	S-TOFHLA ≤ 22	Pictographs (black and white simple line drawings) are effective regardless of race/ethnicity
Choi (2013)	To examine the acceptability and comprehension of instructions in 15 low-literate older adults	United States	Convenience sample N = 15 (82.4% women) Mean age 68.2 years Edu. level: ≤high school (80%) White (73%) S-TOFHLA M = 19.47	Pilot study Cross-sectional study Interviews	Hip replacement surgery	S-TOFHLA ≤ 22	Pictographs are effective for low-literate older adults' discharge instructions in acute health care
Choi (2015)	To examine the effect of pictograph-based discharge instructions on comprehension & recall of older adults with low health literacy	United States	Conv. sample ≥ 65 years w/LHL N = 42 (28 women, 66.7%) Mean age 67.59 years Edu. level: ≤high school (83.3%) White (69%) STOEFLA M = 17.02 (SD = 5.16) Intervention group N = 21 Women (66.7%) White (47.6%) Mean age 68.3 years Edu. level level: ≤high school S-TOFHLA M = 15.95 (SD = 1.43) Comparison group N = 21	RCT 4 weeks discharge instruction education Interviews	Hip replacement surgery	S-TOFHLA ≤ 22	Intervention group had significant improvement in scores on comprehension, $t(35) = -2.29$, $p = 0.02$, and recall, $t(35) = -2.24$, $p = 0.03$, vs. the comparison group

(Continued)

Table 1. (Continued)

Study	Purpose	Setting	Sample characteristics	Design	Disease focus	Literacy measures	Key findings
Chuang et al. (2010)	To compare low-literacy patients and medical staff for preference and comprehension of pictographs intended to illustrate medication use instructions for medical clinic ambulatory patients	South Taiwan	<p>Women (66.7%)</p> <p>White (90.5%)</p> <p>Mean age 66.9 years</p> <p>Edu. level: <high school</p> <p>S-TOFHLA M = 18.1 (SD = 7.09)</p> <p>Convenience sample N = 500</p> <p>Patients n = 250</p> <p>Age M ≤ 60 years (43.2%)</p> <p>Women (54.4%)</p> <p>Men (45.6%)</p> <p>No schooling (61.6%)</p> <p>Medical staff n = 250</p> <p>Age M ≤ 60 years (100%)</p> <p>Women (92%)</p> <p>Male (8%)</p> <p>Edu. level: college degree (86%)</p> <p>Job: Nurse (64.8%)</p>	<p>Prospective study</p> <p>Cross-sectional study</p> <p>Survey</p>	Medication	<p>Education level</p> <p>≤Elementary school</p> <p>(Grades 1–6)</p>	Patients and medical staff had significant differences in preferences for all categories of medical instruction pictographs and had significant differences in comprehension for the pictographs' administration associated with meds. Patients' preferences and comprehension of the medical instruction pictographs were age related
DeWalt et al. (2004)	Development and pilot testing of a disease management program for heart failure patients with low literacy	United States	<p>N = 23</p> <p>Mean age 60 years</p> <p>Men (60%)</p> <p>AA (60%)</p> <p>Income ≤\$15,000/year (74%)</p> <p>Married (67%)</p> <p>Mean education (years): 9.6</p> <p>REALM M: 33% (5th grade)</p> <p>S-TOFHLA: 52% inadequate reading level</p> <p>M knowledge score: 67.3</p> <p>M self-efficacy score (8–32): 30%</p> <p>M MLHFQ score (0–105): 57.3%</p>	<p>Tool development and pilot test</p> <p>Pre-post test</p> <p>Self-report</p> <p>3 months before and after study</p> <p>1 h educational session</p> <p>Supportive phone calls over six weeks</p>	Heart failure	<p>REALM</p> <p>S-TOFHLA</p> <p><9th grade literacy level</p>	Heart failure management program leads to significant improvements in self-care behavior and heart-failure-related symptoms for patients with low literacy

(Continued)

Table 1. (Continued)	Study						
Study	Purpose	Setting	Sample characteristics	Design	Disease focus	Literacy measures	Key findings
Dowse and Ehlers (2001)	To evaluate the use of pharmaceutical pictograms in low-literate South Africans	South Africa	Xhosa ethnic group who had attended school for no more than seven years N = 46 (35 women, 76.1%) Age 41–65 years (56.5%) Edu. level 5–7 years (55.3%)	Pre-post test 3 weeks Interviews	Medication	A short literacy test Education level ≤7 years schooling	Pictographs combined with text are effective for comprehension and recall Pictographs communicate medication information Participants prefer local pictograms ($p < 0.05$)
Dowse and Ehlers (2005)	To determine the influence of medicine labels incorporating pictograms on the understanding of instructions & on adherence	South Africa	Xhosa ethnic group attending an outpatient clinic, who had been prescribed a short course of antibiotics and had attended school for no more than 10 years N = 87 Experimental (Text & pictogram labels) n = 46 (42 women, 91.3%) Age 21–40 years: 45.7% Edu. level 5–7 years (41.3%) Language: isiXhosa (80.4%), English (19.6%) Control (text only) n = 41 (38 women, 92.7%) Age 21–40 years (61%) Edu. level 5–7 years (19, 36.6%) Language: isiXhosa (70.7%), English (34.1%) Percentage for understanding instructions Experimental group: 95% Control group: 70%	RCT Within 3–5 days intervention Interviews	Medication	Education level ≤10 years	Pictographs significantly increase patients' understanding of instructions and adherence ($p < 0.01$) Control group average for understanding, 69.5% Control group average adherence, 71.5%; experimental group, 89.6% ($p < 0.01$)
Kaichman et al. (2013)	To test the efficacy of a pictograph-guided adherence skills building counseling intervention for adults with limited literacy living with HIV	United States	Have HIV & receiving antiretroviral therapy	RCT	HIV	TOFHIA	Interventions or adherence counseling may be required or beneficial for patients with low or marginal literacy skills regardless of their use of pictograms

(Continued)

Table 1. (Continued)

Study	Purpose	Setting	Sample characteristics	Design	Disease focus	Literacy measures	Key findings
Kripalani et al. (2007)	To describe the development, implementation, and preliminary evaluation of an illustrated medication schedule that depicts a patient's daily medication regimen using pill images and icons	United States	<p>N = 446</p> <p>Pictograph guided n = 148</p> <p>Mean age 46.7 years (70% men)</p> <p>Edu. level 12 years</p> <p>HIV + testing: 14 years</p> <p>HIV symptoms: 5.1 years</p> <p>Income < \$10,000/year (69%)</p> <p>TOFHLA < 85% correct (58%)</p> <p>Standard adherence n = 157</p> <p>Mean age 47 years (73% men)</p> <p>Edu. level 12 years</p> <p>HIV + testing: 13.6 years</p> <p>HIV symptoms: 5 years</p> <p>Income < \$10,000/year (71%)</p> <p>TOFHLA < 85% correct (53%)</p> <p>General health improvement</p> <p>n = 141 (65% men)</p> <p>Mean age 47.8 years</p> <p>Edu. level 11.9 years</p> <p>HIV + testing: 13 years</p> <p>HIV symptoms: 4.9 years</p> <p>Income < \$10,000/year (75%)</p> <p>TOFHLA < 85% correct (40%)</p> <p>N = 209</p>	Audio computerized self-interviews	Medication	REALM	Pill card improved medication management and understanding for those with limited health literacy, low levels of education, and cognitive function impairments ($p < 0.05$)
			Frequent use of pill card (83%)	RCT			(Continued)

Table 1. (Continued)

Study	Purpose	Setting	Sample characteristics	Design	Disease focus	Literacy measures	Key findings
Mansoor and Dowse (2003)	To assess the effect of incorporating pictograms on understanding in low-literate participants	South Africa	<p>AA, 191 (91.4%)</p> <p>Women, 122 (58.4%)</p> <p>Mean age 63.7 years</p> <p>Edu. level \geq 12 years (110, 52.6%)</p> <p>REALM: 87 inadequate, 41.6%</p> <p>MMSE \geq 24 (64.6%)</p> <p>All patients had coronary heart disease, 99% had hypertension.</p> <p>Diabetes mellitus: 46%</p> <p>Hypercholesterolemia: 87%</p> <p>Prescribed medications: 6</p> <p>Xhosa low-literacy participants with a maximum of 7 years of formal schooling</p> <p>N = 60 (39 women, 65%)</p> <p>Age 21–40 years (68.3%)</p> <p>Edu. level Grades 5–7th, 40 (66.7%)</p> <p>Experimental group, text and pictogram information</p> <p>n = 30 (60% women)</p> <p>Age 21–40 years (76.7%)</p> <p>Edu. level 5–7th grades (60%)</p> <p>Control, text-only information, asked to read medicine labels and patient information leaflet</p> <p>n = 30 (70% women)</p> <p>Age 21–40 years (60%)</p> <p>Edu. level 5–7th grades (73.3%)</p>	<p>Tool development and intervention</p> <p>3rd month interview</p> <p>RCT</p>	Medication	Education level \leq 7 years schooling	Text-and-pictogram-based information creates a significantly higher level of understanding than text-only information ($p = 0.05$)

Notes: REALM—rapid estimate of adult literacy in medicine; TOFHLA—test of functional health literacy in adults; S-TOFHLA—short test of functional health literacy in adults; AA—African American; Edu.—education; MLHFQ—minnesota living with heart failure questionnaire; MMSE—mini-mental state examination; SEAMS—self-efficacy for appropriate medication use scale.

Table 2. Study characteristics

Study	Research design	N	References*
Descriptive	Cross-sectional	6	Choi (2011, 2012, 2013), Chuang et al. (2010), DeWalt et al. (2004), Kripalani et al. (2007)
Intervention	RCT	5	Choi (2015), Dowse and Ehlers (2005), Kalichman et al. (2013), Kripalani et al. (2007), Mansoor and Dowse (2003)
	Pre-post test	2	DeWalt et al. (2004), Dowse and Ehlers (2001)

*References include duplicate studies.

was no more than high school (Choi, 2012, 2013, 2015). In 3 articles (Dowse & Ehlers, 2001, 2005; Mansoor & Dowse, 2003), the participants' education levels were 5–7 years; in 1 article (Chuang et al., 2010), 61.6% of the participants had no schooling in a patients group and 86% had college degrees in a medical staff group; in another (Kripalani et al., 2007), 52.6% of the participants had more than 12 years of schooling. Seven of the articles (63.6%) (Choi, 2011, 2012, 2013, 2015; DeWalt et al., 2004; Kalichman et al., 2013; Kripalani et al., 2007) presented studies conducted in the US, and the remaining 4 (36.4%) (Chuang et al., 2010; Dowse & Ehlers, 2001, 2005; Mansoor & Dowse, 2003) presented articles conducted elsewhere (South Africa and South Taiwan).

3.2. Studies' characteristics

Five descriptive studies (Choi, 2011, 2012, 2013; DeWalt et al., 2004; Kripalani et al., 2007) consisted of picture-based educational tool development and evaluations using interviews to determine whether a pictographic health education approach was effective or not for people with low health literacy. Of the 7 interventions, 5 (Choi, 2015; Dowse & Ehlers, 2005; Kalichman et al., 2013; Kripalani et al., 2007; Mansoor & Dowse, 2003) were randomized controlled trials; the other 2 (DeWalt et al., 2004; Dowse & Ehlers, 2001) were single-group pre-post studies. The total weighted mean age across the intervention studies was 49.4 years, based on the 3 articles (Choi, 2015; DeWalt et al., 2004; Kripalani et al., 2007) that provided the participants' mean ages. Several different methods were used to collect data: 1 article (Kalichman et al., 2013) used audio-computerized self-interviews (ACASI), and 1 article (DeWalt et al., 2004) used paper self-report data collection. The interventions varied in length from 3 days to 3 months.

3.3. Study designs and picture-based development fields

There were four types of study designs: tool development ($n = 4$), tool development and evaluation by experts ($n = 1$), tool development and evaluation by patients ($n = 3$), and tool evaluation only ($n = 7$) (see Table 3). The health conditions focused upon in the picture-based development were medication adherence ($n = 5$), hip replacement surgery ($n = 3$), breast healthcare ($n = 1$), heart failure ($n = 1$), and human immunodeficiency virus (HIV; $n = 1$).

3.4. Characteristics of pictographs

Studies used pictographs for low health literacy people. The majority of the studies used simple black and white pictographs ($n = 4$). Five studies used pictographs to assist patients with their medications (Chuang et al., 2010; Dowse & Ehlers, 2001, 2005; Kripalani et al., 2007; Mansoor & Dowse, 2003). Most studies used pictographs were conducted in the US ($n = 7$). Ten of 11 articles were the positive effect of increase understanding health education materials.

3.4.1. Health literacy assessment tools

Four different tools were used to assess health literacy: the Test of Functional Health Literacy in Adults (TOFHLA; $n = 1$), the Short Test of Functional Health Literacy in Adults (S-TOFHLA; $n = 4$), the Rapid Estimate of Adult Literacy in Medicine (REALM; $n = 2$), and the Short Literacy Test ($n = 1$), along with education level ($n = 5$).

Table 3. The analysis of study components

Feature	Articles* (N = 11)	References*	Participants** (N = 1,439)
<i>Study design</i>			
Tool development and evaluation by experts	1	Choi (2011)	5
Tool development and evaluation by patients	3	Choi (2012), DeWalt et al. (2004), Kripalani et al. (2007)	238
Only tool evaluation	7	Choi (2013, 2015), Chuang et al. (2010), Dowse and Ehlers (2001, 2005), Kalichman et al. (2013), Mansoor and Dowse (2003)	1,196
<i>Health literacy measurement</i>			
TOFHLA	1	Kalichman et al. (2013)	446
S-TOFHLA	4	Choi (2012, 2013, 2015), DeWalt et al. (2004)	86
REALM	2	DeWalt et al. (2004), Kripalani et al. (2007)	232
Education level	5	Chuang et al. (2010), DeWalt et al. (2004), Dowse and Ehlers (2001, 2005), Mansoor and Dowse (2003)	716
Not present	1	Choi (2011)	5
Short literacy test	1	Dowse and Ehlers (2001)	46
<i>Study setting</i>			
United States	7 (63.6%)	Choi (2011, 2012, 2013, 2015), DeWalt et al. (2004), Kalichman et al. (2013), Kripalani et al. (2007)	746
Other countries	4 (36.4%)	Chuang et al. (2010), Dowse and Ehlers (2001, 2005), Mansoor and Dowse (2003)	693
<i>Study fields</i>			
Hip replacement surgery	3 (27.3%)	Choi (2011, 2013, 2015)	62
Breast healthcare	1 (9.1%)	Choi (2012)	6
Medication	5 (45.4%)	Chuang et al. (2010), Dowse and Ehlers (2001, 2005), Kripalani et al. (2007), Mansoor and Dowse (2003)	902
Heart failure	1 (9.1%)	DeWalt et al. (2004)	23
HIV	1 (9.1%)	Kalichman et al. (2013)	446
<i>Health education materials format</i>			
Only text based	0	None	0
Only pictography based	0	None	0
Simple text and pictography based	11 (100%)	Choi (2011, 2012, 2013, 2015), Chuang et al. (2010), DeWalt et al. (2004), Dowse and Ehlers (2001, 2005), Kalichman et al. (2013), Kripalani et al. (2007), Mansoor and Dowse (2003)	1,439

Notes: TOFHLA—test of functional health literacy in adults; S-TOFHLA—short test.

*References include duplicated studies.

**Number of participants studied, whether participants were experts or patients.

The reading levels in the studies that used the S-TOFHLA were reported as marginal or inadequate (S-TOFHLA \leq 22) in the majority of the studies. If the researchers used years of education, the variables were dichotomized, using the cut off of elementary, less than middle school, or less than high school.

3.5. Relationship between pictography and health education effect

All of the articles described the effectiveness of using pictographs for health education for those with low health literacy. Studies measured effectiveness by using interviews, paper-based self-reports, and questionnaire surveys. Pictography made low health literacy people easy to understand. Ten of the 11 articles (Choi, 2011, 2012, 2013, 2015; Chuang et al., 2010; DeWalt et al., 2004; Dowse & Ehlers, 2001, 2005; Kripalani et al., 2007; Mansoor & Dowse, 2003) found a positive effect of

pictographic educational materials on health self-management, regardless of race/ethnicity and country of origin. However, in Kalichman et al. (2013) the pictograph and adherence to counseling did not improve outcomes in persons with low literacy, though it was effective in persons with moderate health literacy. They surmised that patients with low literacy will need more intense interventions than a brief counseling session to overcome this barrier.

3.6. Effectiveness of pictography for health education

All of the articles (Choi, 2011, 2012, 2013, 2015; Chuang et al., 2010; DeWalt et al., 2004; Dowse & Ehlers, 2001, 2005; Kalichman et al., 2013; Kripalani et al., 2007; Mansoor & Dowse, 2003) combined simple text and pictography. The most common pictographic health education materials featured black and white simple-line drawings with simple text (Choi, 2011, 2012, 2013, 2015). These pictographic health education materials helped to make the information understandable and successfully conveyed healthcare messages to people with health literacy (Choi, 2011, 2012, 2013, 2015). For heart failure patients, the pictographic health education improved self-care behavior (DeWalt et al., 2004). In the medication studies, pictographs with simple text increased the recall of instructions (Chuang et al., 2010), were successful in communicating medication information (Dowse & Ehlers, 2001), were easy to understand (Dowse & Ehlers, 2005; Kripalani et al., 2007; Mansoor & Dowse, 2003) and helped patients remember medication information (Kripalani et al., 2007).

4. Discussion

The 11 articles in this integrative review focused primarily on health literacy measurements, specific health conditions, and the picture-based format of health education materials for people with low health literacy. The review suggests several key findings.

First, communication about health-related subjects with those who have low health literacy is challenging (Arcia et al., 2013). Pictographic representation, however, is beneficial for comprehension and communication in health care (Arcia et al., 2013). Nevertheless, the areas of healthcare that currently use picture-based education for people with low health literacy are limited. In addition, although limited health literacy is affected by age, no studies have focused on the use of pictographic education for the aging population. There is need for more diverse health-related research studies using pictography-based health education materials for elderly.

Currently there are no standard tools for the assessment of health literacy, thus it is difficult to compare levels of health literacy across the studies reviewed here (Berkman et al., 2011). Several of the studies used the dichotomous variable of years of education, but that is not always an appropriate measurement. Health literacy measurements may focus on assessing reading, comprehension, and numeracy (Jordan et al., 2013), but years of education do not provide an accurate measurement of those components. To accurately measure health literacy, future researchers will need more meaningful and consistent health literacy assessment tools.

Finally, picture-based educational materials significantly improved participants' understanding of their health. Most of the studies reported positive results for using pictographic educational materials in healthcare. All 11 of the articles demonstrated the effectiveness of pictographic health education materials increasing patients' learning abilities in different countries, and one article directly pointed out that pictographic health education is effective regardless of race/ethnicity (Choi, 2012). However, one article of HIV education indicated that using picture-based materials did not result in any difference, because people with low health literacy may need more intensive approaches to improving their health. Overall, most of the health education materials currently available are not appropriate for people with a low health literacy level, so improved materials are highly needed.

This integrative review determined what health focus areas were addressed by picture-based health education, the trends that exist in picture-based health education for people with low health literacy, the best health literacy measurements for people with low health literacy, and the

effectiveness of picture-based health education for improving the health status of people with low health literacy.

5. Conclusions

In the US, substantial numbers of adults find it difficult to understand data about their health (Arcia et al., 2013). People's levels of health literacy affect their health and their health management in many ways (U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion, 2010). Low health literacy can lead to greater chances of missing crucial information and therefore causes higher error rates in patients' judgments and decisions about their illnesses (Garcia-Retamero et al., 2012). This results in a direct influence on the status of their illness (Garcia-Retamero et al., 2012). There is a positive relationship between understanding one's health data and improving one's health self-management (Arcia et al., 2013). However, despite attempts to address patients' low health literacy, the majority of materials provided to people with low health literacy are written by people with a high reading level (Badarudeen & Sabharwal, 2008; Goodfellow, Trachimowicz, & Steele, 2008; Helitzer, Hollis, Cotner, & Oestreicher, 2009; Hill-Briggs & Smith, 2008; U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion, 2010) and health-care experts have followed only some of the recommendations made for working with people who have low health literacy (Schwartzberg, Cowett, VanGeest, & Wolf, 2007; U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion, 2010). In this integrative review, we sought to explore which areas of healthcare have used picture-based health education for those with low health literacy and, ultimately, how effective those materials have been. So far, the areas of healthcare using picture-based health education for people with low health literacy have been limited, although the majority of studies indicated a positive effect on learning ability.

5.1. Strengths and limitations

Most studies indicated that using picture-based health education materials for people with low health literacy had a positive effect on health care. But few picture-based health education studies met the inclusion criteria for this integrated review. In addition, the studies were reviewed by only two authors, which may have led to bias; however, the two authors did review the studies three separate times and also subjected their process to review by peer colleagues. There were not many studies that have been done so far; any generalizability of the findings might be limited. Moreover, there were few intervention studies and few participants for assessing the effectiveness of pictographic health care education. In the future, more intervention studies and greater numbers of participants would enable generalizability. In addition, most of the pictographic health care education focused on medication; few studies addressed other healthcare areas. In the future, more research needs to be done on picture-based education in additional areas of healthcare and on a wider range of diseases in order to evaluate its effectiveness.

5.2. Practice implications

In order to address low health literacy needs, healthcare professionals should evaluate their current patient education for reading level and low-literacy acceptability. Reading levels can be assessed using the health measurements tools such as S-TOFHLA, and REALM. Patient education handouts should be on a reading level below 5th-grade level (Kirsch & Educational Testing Service, & National Center for Education Statistics, 1993). If there is a lack patient education materials tailored for low health literacy, healthcare professionals can create patient education material using picture-based information. Pictures should be simple, clear, and culturally tailored for easy to understand. Appropriate picture-based health education materials would make it easier for such people to understand health information, as opposed to text-based health education materials alone.

Acknowledgements

I wish to take this opportunity to express my deep regard to Dr Sharon A. Brown, PhD, RN, FAAN, for imparting her expert knowledge about literature reviews and guiding me in this research. Editorial support with manuscript development was provided by the Cain Center for Nursing Research and the Center for Transdisciplinary Collaborative Research in Self-management Science (P30, NR015335) at The University of Texas at Austin, School of Nursing.

Funding

The authors received no direct funding for this research.

Competing Interests

The authors declare no competing interests.

Author details

Jungmin Park¹

E-mail: jungminpark@utexas.edu

Julie Zuniga¹

E-mail: jzuniga@nursing.utexas.edu

¹ School of Nursing, The University of Texas at Austin, 1700 Red River St., Austin, TX 78701, USA.

Citation information

Cite this article as: Effectiveness of using picture-based health education for people with low health literacy: An integrative review, Jungmin Park & Julie Zuniga, *Cogent Medicine* (2016), 3: 1264679.

References

- Arcia, A., Bales, M. E., & William Brown III, M. C. (2013). Method for the development of data visualizations for community members with varying levels of health literacy. *AMIA Annual Symposium Proceedings*, 2013, 51–60.
- Austin, P. E., Matlack, R., Dunn, K. A., Kesler, C., & Brown, C. K. (1995). Discharge instructions: Do illustrations help our patients understand them? *Annals of Emergency Medicine*, 25, 317–320. [http://dx.doi.org/10.1016/S0196-0644\(95\)70286-5](http://dx.doi.org/10.1016/S0196-0644(95)70286-5)
- Badarudeen, S., & Sabharwal, S. (2008). Readability of patient education materials from the American academy of Orthopaedic surgeons and pediatric orthopaedic society of north America web sites. *The Journal of Bone and Joint Surgery-American Volume*, 90, 199–204. <http://dx.doi.org/10.2106/JBJS.G.00347>
- Berkman, N. D., Sheridan, S. L., Donahue, K. E., Halpern, D. J., & Crotty, K. (2011). Low health literacy and health outcomes: An updated systematic review. *Annals of Internal Medicine*, 155, 97–107. doi:10.7326/0003-4819-155-2-201107190-00005
- Choi, J. (2011). Pictograph-based discharge instructions for low-literate older adults after hip replacement surgery: Development and validation. *Journal of Gerontological Nursing*, 37, 47–56. doi:10.3928/00989134-20110706-03
- Choi, J. (2012). Development and pilot test of pictograph-enhanced breast health-care instructions for community-residing immigrant women. *International Journal of Nursing Practice*, 18, 373–378. doi:10.1111/j.1440-172X.2012.02051.x
- Choi, J. (2013). Older adults' perceptions of pictograph-based discharge instructions after hip replacement surgery. *Journal of Gerontological Nursing*, 39, 48–54. doi:10.3928/00989134-20130415-02
- Choi, J. (2015). *Effect of pictograph-based discharge instructions on older adults' comprehension and recall: A pilot Study. Research in Gerontological Nursing*, 9, 66–71. doi:10.3928/19404921-20150513-05
- Chuang, M.-H., Lin, C.-L., Wang, Y.-F., & Cham, T.-M. (2010). Development of pictographs depicting medication use instructions for low-literacy medical clinic ambulatory patients. *Journal of Managed Care Pharmacy*, 16, 337–345. <http://dx.doi.org/10.18553/jmcp.2010.16.5.337>
- Davis, T. C., Long, S. W., Jackson, R. H., Mayeaux, E., George, R. B., Murphy, P. W., & Crouch, M. A. (1993). Rapid estimate of adult literacy in medicine: A shortened screening instrument. *Family Medicine*, 25, 391–395.
- Delp, C., & Jones, J. (1996). Communicating information to patients: The use of cartoon illustrations to improve comprehension of instructions. *Academic Emergency Medicine*, 3, 264–270. doi:10.1111/j.1553-2712.1996.tb03431.x
- DeWalt, D. A., Pignone, M., Malone, R., Rawls, C., Kosnar, M. C., George, G., ... Angel, B. (2004). Development and pilot testing of a disease management program for low literacy patients with heart failure. *Patient Education and Counseling*, 55, 78–86. doi:10.1016/j.pec.2003.06.002
- Doak, C. C., Doak, L. G., Friedell, G. H., & Meade, C. D. (1998). Improving comprehension for cancer patients with low literacy skills: Strategies for clinicians. *CA: A Cancer Journal for Clinicians*, 48, 151–162. doi:10.3322/canjclin.48.3.151
- Dowse, R., & Ehlers, M. (2001). The evaluation of pharmaceutical pictograms in a low-literate South African population. *Patient Education and Counseling*, 45, 87–99. doi:10.1016/S0738-3991(00)00197-X
- Dowse, R., & Ehlers, M. (2005). Medicine labels incorporating pictograms: Do they influence understanding and adherence? *Patient Education and Counseling*, 58, 63–70. doi:10.1016/j.pec.2004.06.012
- Garcia-Retamero, R., Okan, Y., & Cokely, E. T. (2012). Using visual aids to improve communication of risks about health: A review. *The Scientific World Journal*, 2012, 562637. doi:10.1100/2012/562637
- Goodfellow, G. W., Trachimowicz, R., & Steele, G. (2008). Patient literacy levels within an inner-city optometry clinic. *Optometry-Journal of the American Optometric Association*, 79, 98–103. doi:10.1016/j.optm.2007.03.015
- Helitzer, D., Hollis, C., Cotner, J., & Oestreicher, N. (2009). Health literacy demands of written health information materials: An assessment of cervical cancer prevention materials. *Cancer Control*, 16, 70–78.
- Hill-Briggs, F., & Smith, A. S. (2008). Evaluation of diabetes and cardiovascular disease print patient education materials for use with low-health literate populations. *Diabetes Care*, 31, 667–671. doi:10.2337/dc07-1365
- Jordan, J. E., Buchbinder, R., Briggs, A. M., Elsworth, G. R., Busija, L., Batterham, R., & Osborne, R. H. (2013). The Health Literacy Management Scale (HeLMS): A measure of an individual's capacity to seek, understand and use health information within the healthcare setting. *Patient Education and Counseling*, 91, 228–235. <http://dx.doi.org/10.1016/j.pec.2013.01.013>
- Kalichman, S. C., Cherry, C., Kalichman, M. O., Amaral, C., White, D., Grebler, T., ... Schinazi, R. F. (2013). Randomized clinical trial of HIV treatment adherence counseling interventions for people living with HIV and limited health literacy. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 63, 42–50. doi:10.1097/QAI.0b013e318286ce49
- Kirsch, I. S., Educational Testing Service, & National Center for Education Statistics. (1993). *Adult literacy in America: A first look at the results of the national adult literacy survey*. Washington, DC: Office of Educational Research and Improvement, US Dept. of Education.
- Kripalani, S., Robertson, R., Love-Ghaffari, M. H., Henderson, L. E., Praska, J., Strawder, A., ... Jacobson, T. A. (2007). Development of an illustrated medication schedule as a low-literacy patient education tool. *Patient Education and Counseling*, 66, 368–377. doi:10.1016/j.pec.2007.01.020
- Kutner, M., Greenburg, E., Jin, Y., & Paulsen, C. (2006). *The health literacy of America's adults: Results from the 2003 national assessment of adult literacy*. NCEES 2006-483. Washington, DC: National Center for Education Statistics.

- Mansoor, L. E., & Dowse, R. (2003). Effect of pictograms on readability of patient information materials. *The Annals of Pharmacotherapy*, 37, 1003–1009. doi:10.1345/aph.1C449
- Mayer, R. E. (1947/2014). *The Cambridge handbook of multimedia learning* (pp. 43–71, 2nd ed.). New York, NY: Cambridge University Press.
- Mayer, R. E., & Ebooks Corporation. (2009). *Multimedia learning* (2nd ed.). Leiden: Cambridge University Press.
<http://dx.doi.org/10.1017/CBO9780511811678>
- Nielsen-Bohlman, L., NetLibrary, I., & Institute of Medicine (US) Committee on Health Literacy. (2004). *Health literacy: A prescription to end confusion*. Washington, DC: National Academies Press.
- Paasche-Orlow, M. K., Parker, R. M., Gazmararian, J. A., Nielsen-Bohlman, L. T., & Rudd, R. R. (2005). The prevalence of limited health literacy. *Journal of General Internal Medicine*, 20, 175–184.
doi:10.1111/j.1525-1497.2005.40245.x
- Schwartzberg, J. G., Cowett, A., VanGeest, J., & Wolf, M. S. (2007). Communication techniques for patients with low health literacy: A survey of physicians, nurses, and pharmacists. *American Journal of Health Behavior*, 31, 96–104.
<http://dx.doi.org/10.5993/AJHB.31.s1.12>
- Sorden, S. D. (2012). The cognitive theory of multimedia learning. In B. Irby, G. H. Brown, R. Lara-Alecio, & S. A. Jackson (Eds.), *Handbook of educational theories* (p. 1). Charlotte, NC: Information Age Publishing.
- U.S. Department of Health and Human Services, & Office of Disease Prevention and Health Promotion (2010). *National action plan to improve health literacy*. Washington, DC.



© 2016 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format
Adapt — remix, transform, and build upon the material for any purpose, even commercially.
The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.
You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



Cogent Medicine (ISSN: 2331-205X) is published by Cogent OA, part of Taylor & Francis Group.

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

