



Received: 02 September 2016
Accepted: 02 January 2017
First Published: 06 January 2017

*Corresponding author: James A. Levine, Health Solutions, Mayo Clinic Arizona and Arizona State University, Suite 200 CRB, 13400 East Shea Blvd, Scottsdale, AZ 85259, USA
E-mail: levine.james@mayo.edu

Reviewing editor:
Udo Schumacher, University Medical Center Hamburg-Eppendorf, Germany

Additional information is available at the end of the article

PSYCHIATRY | REVIEW ARTICLE

Mental health issues in survivors of sex trafficking

James A. Levine^{1*}

Abstract: There are more than three million trafficked women, many of who are children. This population has been largely forgotten because the victims are geographically transitory and exist within an illegal framework. Research, often conducted using invalid techniques, suggests that most survivors of sex trafficking (i.e. those evaluated in shelters), have mental illness. The majority of survivors have depression, anxiety, posttraumatic stress disorder or a more severe diagnosis: Disorders of Extreme Stress (DESNOS). In addition to these diagnoses, many victims of sex trafficking also have secondary psychological issues such as alcohol and drug abuse plus concurrent medical illnesses, which add to the psychological burden that sex trafficked victims endure. Mental health interventions often focus on identifying potential victims in healthcare centers and public places such as shopping malls or at truck stops. The problem, however, is that once a sex trafficked person is rescued there are no prospective clinical trials to guide therapy; oftentimes by default, trauma-based cognitive behavioral therapy is used; such approaches may not be effective for those with DESNOS. Considering the number of women and children who have been victimized by sex trafficking, it is astonishing that so little is known about the natural history of mental health issues and almost nothing is known about how best to treat these victims.

Subjects: Public Health Policy and Practice; Social Work and Social Policy; Pediatrics & Child Health; Psychiatry; Specialist Community Public Health Nursing

Keywords: human sex trafficking; children; mental health; social determinants of health

1. Introduction

There are approximately 3.3 million sex-trafficked persons in the world, nearly all of them women (ILO Global Estimate of Forced Labour: results & methodology/International Labour Office, Special Action Programme to Combat Forced Labour (SAP-FL), 2012). Although data from 61 countries



James A. Levine

ABOUT THE AUTHOR

James A. Levine is professor of Health Solutions and Director of Obesity Solutions at Mayo Clinic and Arizona State University. He has worked with the United Nations to prevent child exploitation with numerous organisations in India, Africa and the United States. He is on the board of the International Center for Missing and Exploited Children. His novel *The Blue Notebook* was published in 26 countries and 17 languages focused on child sex trafficking.

PUBLIC INTEREST STATEMENT

Survivors of sex trafficking bare many scars of their exploitation; some are physical, many are mental. In this paper the current knowledge is reviewed regarding the mental health implications of sex trafficking. The causality, pathophysiology and current treatment options are discussed and suggestions are made to improve the mental health apparatus and research agenda for victims of sex trafficking.

suggest that the ratio of women to girls (a girl being less than 18 years of age) is 5:1, many agencies believe that child sex trafficking far exceeds one million children (United Nations Office on Drugs & Crime, 2009). The precise numbers of woman being trafficked for the purpose of sexual exploitation (“sex trafficked”) is not accurately known because the activity is criminal in about 80% of countries and so the quality of information is poor. In this paper the implications of sex trafficking on mental health is examined, the limitations of current knowledge are explained and priorities for future research are identified.

2. Causes of mental health issues in sex trafficking


Sex trafficking and mental illness seem inevitably linked. However, understanding why sex trafficking is associated with mental illness helps better understand the depth of the victimization and how better to plan for comprehensive therapy.

2.1. Biological factors

The high levels of psychological stress associated with sexual exploitation are associated with neurochemical and structural changes in the brain; more so because many sex-trafficked victims are children. It can be conjectured that the levels of trauma associated with sex trafficking cause changes in hypothalamic nuclei gene expression and altered release of, pro-opiomelanocortin, corticotropin-releasing hormone, adrenocorticotrophic and cortisol, a key stress hormone (Busso, McLaughlin, & Sheridan, 2016; Hewagalamulage, Clarke, Rao, & Henry, 2016; Mills, Carter, Rudd, Claxton, & O'Brien, 2016; Mogami et al., 2016; Osterlund et al., 2016; Provenzi et al., 2016). Intense psychological stress impacts other neuromodulator systems too such as opioids (Burke, Finn, McGuire, & Roche, 2016; Szklarczyk et al., 2016), GABA (Liu et al., 2016; Yang et al., 2016), norepinephrine (Thakare, Dhakane, & Patel, 2016; Wood, Valentino, & Wood, 2016) 5-HT (Demirdas, Naziroglu, & Unal, 2016; Kumar, Rajkumar, Lee, & Dawe, 2016; Li-Tempel et al., 2016) and even growth hormone (Aggarwal et al., 2014; Marina, Klose, Nordenbo, Liebach, & Feldt-Rasmussen, 2015; Savendahl, 2012). These factors, shown in animal models, subsequently impair how the body handles stress (Burke et al., 2016; Demirdas et al., 2016; Kumar et al., 2016; Li-Tempel et al., 2016; Szklarczyk et al., 2016).

The structure of the brain changes with trauma too. Environmental exposure directly affects neuronal formation and brain anatomy (Corr & Perkins, 2006; Windholz, 1987). Multiple neuroplasticity factors control brain remodeling in response to environmental change (Diaz Heijtz & Forssberg, 2015; Simchon Tenenbaum, Weizman, & Rehavi, 2015). Substantial stress is directly associated with neuroplasticity-associated changes in brain structure and function (McEwen, Gray, & Nasca, 2015; Vyas et al., 2016; Wilson, Grillo, Fadel, & Reagan, 2015). A vicious cycle exists (Figure 1) whereby

Figure 1. Cartoon to show relationship between chronic stress and changes in brain biology.



chronic stress impacts brain structure and function rendering the growing brain more vulnerable to the ill effects of abuse.


2.2. Genetic factors

Many sex trafficked victims, like many victims of violence, are from families with high prevalence of depression (Xia & Yao, 2015), schizophrenia (Chiapponi, Piras, Piras, Caltagirone, & Spalletta, 2016; Davis et al., 2016), alcoholism (Matošić, Marusic, Vidrih, Kovak-Mufic, & Cicin-Sain, 2016) and/or, substance/drug abuse (Yu & McClellan, 2016). No specific studies have been conducted on genetic linkages and mental health disorders in these disorders, however. Each of these disorders can be associated with genetic factors and so survivors of sex trafficking may be genetically predisposed to develop mental health issues (Ehlert, 2013; Serafini et al., 2015). Also, genetics factors influence how a person responds to stress and to the risk of developing Posttraumatic Stress Disorder (Lebois, Wolff, & Ressler, 2016). Epigenetic influences are *in utero* exposures that impact fetal development (Anacker, O'Donnell, & Meaney, 2014; Lutz, Almeida, Fiori, & Turecki, 2015; Provençal & Binder, 2015). Maternal alcohol (Gupta, Gupta, & Shirasaka, 2016) and illicit drug use (Antonelli, Pallares, Ceccatelli, & Spulber, 2016) and even *in utero* violence against the mother (Anacker et al., 2014; Lutz et al., 2015; Provençal & Binder, 2015) are examples of factors that can impair the mental status of the child. Thus both genetic and epigenetic factors may predispose sex-trafficked individuals to develop mental health issues.

2.3. Psychosocial factors

Sex trafficking victims are immersed in a social ecology that is linked with other deleterious factors such as crime, drug abuse and poverty (Banovic & Bjelajac, 2012). This circle of violence (Figure 2), independently has a negative impact on mental health (Diehl, Pillon, dos Santos, Rassool, & Laranjeira, 2016; Huston, Anglin, & Eckstein, 1996; Kimerling, Alvarez, Pavao, Kaminski, & Baumrind, 2007).

Figure 2. Social factors associated with mental illness in sex trafficking.



3. Prevalence of mental health issues in sex trafficking

Research on the prevalence of mental health issues associated with sex trafficking focuses on survivors. There are no studies on the incidence of mental health issues in people actively being trafficked, because identifying new cases of mental illness while a person is being sexually exploited breaches the ethical obligation to intervene.

3.1. Prevalence data

Prevalence data are inconsistent in this population because of, (a) the different sex trafficked populations that have been studied, (b) the differing definitions of mental illness and (c) the variability in the quality of work. In one example, Abas et al. (2013) examined a historical cohort of adult sex trafficked survivors who had returned to Moldova after registering for assistance with an international migration agency. Out of 176 survivors, data from 120 women were examined. DSM-IV Criteria (Frances, Pincus, Widiger, Davis, & First, 1990) were used to categorize mental disorders. Fifty-four percent had mental illness as defined by the criteria; 36% had PTSD, 13% had depression without PTSD and 6% had another anxiety disorder. In contrast, Crawford examined a cohort of 80 Nepalese women aged 12–19 who were survivors of sex trafficking (Crawford & Kaufman, 2008). They identified, through detailed examination of case files, that somatic symptoms outweighed behavioral ones. In this study, somatic symptoms of headache, itching, social withdrawal, aggression, stomach pain, fatigue, altered behaviors towards males, pelvic pain and low motivation were the principal mental health sequelae of sex trafficking. The comparison of these two studies highlights common issues regarding mental health research in sex trafficking survivors.

Oram, Stöckl, Busza, Howard and Zimmerman (2012) collated data from four prevalence studies (Cwikel, Chudakov, Paikin, Agmon, & Belmaker, 2004; Hossain, Zimmerman, Abas, Light, & Watts, 2010; Ostrovschi et al., 2011; Tsutsumi, Izutsu, Poudyal, Kato, & Marui, 2008) of sex trafficked survivors. The methodologies included checklists, psychological scales, and clinical interviews. Overall, 78% of survivors experience clinical anxiety (range 48–98%); 52% depression (range 3–100%) and 37% posttraumatic stress disorder (range 8–77%). Beyond formal diagnoses of mental illnesses, survivors of sex-trafficking report many other symptoms (Ijadi-Maghsoodi, Cook, Barnert, Gaboian, & Bath, 2016; Shandro et al., 2016) such as feelings of fear and distrust, being trapped, hopelessness, shame, humiliation, a sense of continuous stress, anger, and irritability. Survivors also often report poor quality sleep, insomnia, and nightmares. Survivors of sex trafficking often feel socially stigmatized, whereby they cannot return to their homes and communities because they will be pre-identified as being sex workers. In these instances, sex trafficking survivors can report a sense of being re-victimized even after rescue (Ijadi-Maghsoodi et al., 2016).

Overall, the vast majority of survivors of sex trafficking have mental illness especially anxiety disorder, depression or, posttraumatic stress disorder. There are other mental health sequelae too such as poor sleep, social stigmatization and feelings such as those of hopelessness.

3.2. Physical illness secondary to mental trauma in sex trafficking survivors

There are physical consequences associated with mental illness in survivors of sex trafficking (Crawford & Kaufman, 2008; Cwikel et al., 2004; Ehlert, 2013; Hossain et al., 2010; Oram, Ostrovschi et al., 2012; Richards, 2014; Schaefer et al., 2015; Shandro et al., 2016; Zimmerman, Hossain, & Watts, 2011; Zimmerman et al., 2008). The skin and hair may show evidence of cigarette burns, attempts at self-harm, rashes, evidence of intravenous drug use, vitamin deficiency, bruising from physical abuse and infectious diseases such as tuberculosis or lice. The teeth need to be examined for dental caries and poor overall dental hygiene; methamphetamine and bulimia scar teeth too. Although survivors of sex trafficking can be malnourished, obesity may be present and this should not dissuade a clinician from concern. Gynecological issues include evidence of sexually transmitted diseases (STD's), cervical dysplasia, non-menstrual vaginal bleeding, vaginal pain, dysuria, dyspareunia, traumatic scarring and ovulatory failure. The cardiovascular system can be impacted by hypertension and ischemic disease. In addition, survivors of sex trafficking with mental illness may have concomitant chronic pain syndromes, dizziness, fainting and fibromyalgia. X-rays may show prior

abuse-related fractures and decreased bone density. Immune function may be impaired not only because of HIV but also because of malnutrition. Hepatitis B and C need to be screened for. Survivors of sex trafficking who are predisposed to asthma or diabetes are inadequately cared for.

3.3. Secondary social sequelae

Survivors of sex trafficking with mental illness may have attendant drug, cigarette and alcohol abuse (Chung & English, 2015; Davidson, Grigorenko, Boivin, Rapa, & Stein, 2015; Gibbons & Stoklosa, 2016). Rehabilitated survivors of sex trafficking are often stigmatized, prone to become victims of violence, pressured to not use barrier contraception (Decker, Mack, Barrows, & Silverman, 2009; Mantini, 2008) and are more likely to become involved in criminality (Diehl et al., 2016; English, 2015; Greenbaum, Dodd, & McCracken, 2015; Houston-Kolnik, Todd, & Wilson, 2016). In some communities survivors, who are viewed as tainted, are unable to marry (Rimal & Papadopoulos, 2016). These types of social issues need to be considered when providing mental health services to survivors.

4. Mental health treatment in survivors of sex trafficking

There are three principal areas of focus; (1) early detection of sex trafficked victims, (2) identification of the most effective mental health treatment and (3) organization of multi-disciplinary healthcare teams.

4.1. Early detection of sex trafficked victims so that they can be identified, protected and treated for mental health issues

The Polaris Project identified a list of risk factors that a person is at risk of being sex trafficked (Figure 3) (Committee opinion No. 507: Human trafficking, 2011; Helton, 2016; Shandro et al., 2016). Although, they are well accepted there are no published data on the criteria's positive predictive value (correctly rule-in disease) or negative predictive value (correctly rule out disease). Nonetheless these criteria have been used to help identify potential victims in number of settings such as with truck drivers, police and hoteliers (Institute of Medicine & National Research Council, 2014; Jana, Dey, Reza-Paul, & Steen, 2014; Roe-Sepowitz, Gallagher, Risinger, & Hickle, 2015). In the United States there is a national telephone hotline run by the National Human Trafficking Resource Center [<https://traffickingresourcecenter.org>] on 1-888-373-7888 where advice can be obtained.

Figure 3. Polaris Project (3). General indicators that a person may be at risk of being sex-trafficked.

- Age < 18 years old
- Multiple sexual partners
- Multiple sexually transmitted diseases
- Inappropriate attire [e.g. lingerie]
- Tattoos or branding
- Sexual abuse, genital trauma
- Lack of official identification
- Vague answers to questions
- Inconsistencies in story
- No eye contact
- No control of money [some one else controls]
- Malnourishment
- Signs of physical abuse:
 - burns
 - bruises
 - broken bones
- Signs of Depression or, Post Traumatic Stress Disorder
- Drug/alcohol addiction

Health care facilities are especially important for identifying potential victims of sex trafficking (Grace et al., 2014; Nuzzolese, 2014; Ross et al., 2015) because medical assistance is often necessary to keep a sexually exploited person working. Greenbaum et al. (2015) used cross-sectional emergency room data to identify six key questions to identify a person who is at risk of being sex-trafficked;

- (1) Is there a previous history of drug and/or alcohol use?
- (2) Has the youth ever run away from home?
- (3) Has the youth ever been involved with law enforcement?
- (4) Has the youth ever broken a bone, had traumatic loss of consciousness, or sustained a significant wound?
- (5) Has the youth ever had a sexually transmitted infection?
- (6) Does the youth have a history of sexual activity with more than 5 partners?

When the patient has four or more positive responses, the positive predictive value for being sexually exploited is, 88% and the negative predictive value, 88% (Greenbaum et al., 2015). There are other indicators in a healthcare facility that a person may be being sex trafficked. For example, sex trafficking victims are often accompanied by a female accomplice of the pimp called a, “Bottom Girl”. This is to ensure that she does not attempt escape; the Bottom Girl will control the money and papers of the victim. Also sex trafficked victims can be vulnerable to Stockholm syndrome when they identify romantically with their pimp (Chesnay, 2013).

If a healthcare provider identifies that a patient is being sex trafficked it is likely that the patient has mental illness. The following general recommendations facilitate the immediate interaction with the patient; (a) The patient needs to be separated from an accompanying person (e.g. Bottom Girl or, pimp), (b) it is important to talk directly to the victim, if necessary using an interpreter, (c) questions should be asked to ascertain whether the woman/girl is independent (e.g. has self control over her money and identity papers), (d) it is important to immediately assess the patient’s mental status, particularly with risk for self-harm, in order to ascertain the urgency for intervention, (e) protective services and law enforcement should be contacted.

4.1.1. Short-term intervention

Once a victim has been identified, a mental health plan needs to be formulated. The communication style with the patient must be open, trustworthy, safe, reliable, confidential, consistent, reassuring, knowledgeable, and non-judgmental (Chesnay, 2013). Early mental health intervention is important as psychiatric issues can be life threatening (e.g. suicidal ideation). Underlying psychiatric disorders, alcohol and drug abuse need to be treated. In general, medications should be used only for specific psychiatric indications. Benzodiazepines are especially discouraged as they can exacerbate dissociation (Ratzer, Brink, Knudsen, & Elklit, 2014) although there have not been randomized pharmaceutical trials in this patient group. General health evaluation is necessary too including gynecological examinations and screening for sexually transmitted diseases (Oram, Ostrovski et al., 2012). The legal (e.g. guardianship) and immigration status of the patient must be established; specific visas exist to protect sex trafficked persons. Social workers are critical to help identify resources and identify safe temporary housing most often in protected shelters or mental health facilities.

Health professionals treating survivors of sex trafficking can follow a general set of mental health principles (Chesnay, 2013). It is important to, (A) Acknowledge the trauma. (B) Explore the trauma at an acceptable pace to the survivor. (C) Explore the person’s self-identity to avoid re-victimized nation and stigmatization. (D) Participate in the survivor setting new life goals and helping individual plan to meet these goals. (E) Set in place not only continued mental health support but also general health surveillance. (F) Encourage good quality sleep, physical activity and nutrition. (G) Help the patient reframe experiences to identify positive attributes of the individual and recognize their

accomplishments. (H) Establish support for life skills such as financial intelligence, healthy recreation and maintaining healthy relationships. Once the patient is stabilized medically, legally and socially, a longer-term treatment plan can be defined.

4.1.2. Long-term intervention

There are no clinical trials that identify the best modality of mental health treatment for survivors of sex trafficking; trauma-based cognitive behavioral therapy is one of the most commonly used treatment (Cohen, 2008; Keeshin & Strawn, 2014; Walter, Buckley, Simpson, & Chard, 2014). Trauma focused cognitive behavioral therapy uses 12–20 structured sessions to teach the patient coping skills, trauma narration, trauma processing, treatment consolidation and closure.

There is an important area of emerging research that suggests that PTSD (a concept originally developed to explain the complex mental health challenges of veterans returning from Vietnam) is inadequate for defining the chronic and repeated traumatic exposures that survivors of sex trafficking have experienced (van der Kolk, 2001). The conglomerate of symptoms that sex trafficked survivors experience may fit under a newer descriptor known as; Disorders of Extreme Stress (DESNOS) or Complex PTSD (Hyland et al., 2016; Karatzias et al., 2016; Sachser, Keller, & Goldbeck, 2016). This is important because this diagnosis impacts outcome (Ford & Kidd, 1998) and may be a consideration for designing effective treatments for survivors of sex trafficking. Extensive work has been conducted on how best to treat DESNOS (van der Kolk, 2001) but this knowledge has not been adapted and trialed in survivors of sex-trafficking.

A range of other supportive/alternative therapies have been used too (Recommendations to reduce psychological harm from traumatic events among children and adolescents, 2008); for example, art therapy (Wong, 2008), Narrative Exposure Therapy (Kangaslampi, Garoff, & Peltonen, 2015; Larsen, Wiltsey Stirman, Smith, & Resick, 2016), music therapy (Froehlich, 1984), equine therapy (del Rosario-Montejo, Molina-Rueda, Muñoz-Lasa, & Alguacil-Diego, 2015) and eye movement desensitization and reprocessing therapy (Acarturk et al., 2016; Nijdam & Olff, 2016). The latter, for example, is especially of interest in treating DESNOS (van der Kolk, 2001). Despite the conceptual attractiveness of many of these approaches there have not been any clinical trials of efficacy or effectiveness.

5. Healthcare organization to support victims of sex trafficking

Individual healthcare facilities need a plan to identify and manage victims of sex trafficking. It is important to, (a) establish and list key contacts with local police forces and other related agencies such as the FBI, INS and National Center for Missing and Exploited Children; (b) victims need to be guaranteed safety whilst in the healthcare facility and so physical security, surveillance cameras, and door locks need to be reviewed; (c) a protocol needs to be in place to establish the legal status of the victim; (d) a communications plan is required; for example, preprogramming telephones with emergency numbers and the number for the national hotline (888-373-7888), plus a communications policy needs to be established to cover communication with a victim's family and protect the victim from contact by pimps.

The mental/social care plan for survivors must be designed for long-term sustainability because recidivism rates are high. Child welfare agencies are legally obliged under the Preventing Sex Trafficking and Strengthening Families Act, P.L. 113–183, (US Congress, 2014)

To develop policies and procedures to identify, document, and determine appropriate services for children under the placement, care, or supervision of a child welfare agency and who are at risk of becoming sex trafficking victims or who are victims of sex trafficking.

Multidisciplinary teams are needed to include psychiatric, medical, law enforcement and social work professionals. For example, the mental health care of sex trafficked survivors is enhanced in states such as Minnesota that have Safe Harbor legislation; whereby children who are caught by law

enforcement engaged in prostitution cannot be charged with a related crime (Minnesota Department of Health, 2014). Lastly, many staff who work extensively with sex-trafficked survivors feel burned out and so plans are needed to support the psychological needs of healthcare staff (Kliner & Stroud, 2012).

6. Mental health issues in sex trafficking: Limitations to current knowledge

High quality research is important in building the case for policy change, the allocation of healthcare resources and the funding necessary to support the social services that sex-trafficked survivors need (Cary, Oram, Howard, Trevillion, & Byford, 2016). There are several reasons why research is so limited in understanding mental health issues associated with sex trafficking.

6.1. The concepts of sex trafficking are grossly oversimplified

Mental health issues in sex trafficked victims often are multiple; e.g. trauma *plus*, sexual abuse, physical abuse, kidnapping, social disadvantage, educational curtailment and drug abuse. Conceptualizing sex trafficking as an isolated issue oversimplifies a multi-layered problem.

6.2. Research is often conducted on unrepresentative samples

Mental health research includes only survivors of sex trafficking who are living in rescue centers, which is an unrepresentative sample. Also, research in this area is most often conducted by agencies with an inherent interest in the outcomes of the research (e.g. to gain funding).

6.3. Missing data

Many victims do not actually know their age, heritage or family details. The magnitude of the trauma often results in subjects being unable to complete the research protocol.

6.4. Many of the research tools have not been validated in sex trafficked victims (Cannon, Arcara, Graham, & Macy, 2016)

Sex trafficked victims may not be conversant in the language where the study is being conducted nor with cultural norms. As noted above, the multilayered nature of the trauma sex trafficked victims experience necessitates the need for correctly validated instruments.

6.5. Ethical issues

People who survive sex trafficking are already victims of coercion. Thus, obtaining informed consent is especially complex. When a health worker asks a survivor to participate in a research project, it is difficult to avoid a sense of implicit coercion.

6.6. Does research re-stigmatize victims?

The nature of research may be re-stigmatizing. For example, a consent form may have the title, "Mental health issues in sex trafficked victims." This re-victimizes the survivor to identify herself in this fashion.

6.7. Low priority research area

Clinicaltrial.gov reports that there are currently 55,610 active clinical trial in Cancer and 9,505 trials in depression. There are two clinical trials in sex trafficking. By funding agencies consistently not supporting research in this area, the failure to gain useful knowledge and develop effective treatments is perpetuated.

6.8. Policy and impact

New policies to protect victims of sex trafficking have unknown impact.

7. Mental health issues in sex trafficking: Recommendations

Recommendation 1. *Techniques*. International standards on how mental illness is defined in these individuals are necessary. Valid techniques are needed to assess the allostatic load (psychological plus psychological stressors) that sex trafficked victims endure (McEwen & Wingfield, 2003).

Recommendation 2. *Natural history*. Better information is needed about the nature, causal factors and natural history of mental illness in survivors of sex-trafficking. The course of mental illness in sex-trafficked victims is unknown. Most survivors of sex trafficking have mental illness, which is associated with multiple exacerbating and perpetuating factors. Little is understood on how these affect the long-term outcomes of survivors of sex trafficking compared to other patients. Understanding long-term outcomes would help define the resources that are needed at rescue.

Recommendation 3. *Treatment*. Clinical trials are needed to understand how best to treat mental illness in sex trafficked victims.

Recommendation 4. *Research ethics*. In research on victims who are currently being exploited, there will need to be a declaration of intent to withdraw the research subject from sex trafficking. It is important to avoid, in the research process, re-stigmatizing a person as a victim of sex trafficking/prostitution. Caution is necessary to avoid coercing survivors to engage in consent.

Recommendation 5. *Funding*. There are more than three million sex trafficked persons; many of whom are young. Little is known about the psychological and physical burden they endure. Nothing is known about the best method of intervention. The size of the population and its youth justify allocating research resources in this domain.

Recommendation 6. *Health care systems research*. A guide that includes protocols and procedures is needed to enable health care systems to best help potential victims of sex trafficking who they encounter. We need to establish national data management systems that track the magnitude of the problem, resource management and the effectiveness of interventions.

8. Conclusion

Most of the three million women and girls who are currently being sex trafficked have mental illness. Considering the size of the problem and magnitude of the trauma so little is known about the cumulative misery these young people endure and what to do about it. That needs to change.

Funding

This work was funded by NIH grants, R42 DK085738-02, 5R01DK085516-05 and R015R01HL114024-1 and a CODE Award from Mayo Clinic Center for Innovation.

Competing Interests

The author declares no competing interest.

Author details

James A. Levine¹
E-mail: levine.james@mayo.edu

¹ Health Solutions, Mayo Clinic Arizona and Arizona State University, Suite 200 CRB, 13400 East Shea Blvd, Scottsdale, AZ 85259, USA.

Citation information

Cite this article as: Mental health issues in survivors of sex trafficking, James A. Levine, *Cogent Medicine* (2017), 4: 1278841.

References

- Abas, M., Ostrovski, N. V., Prince, M., Gorceag, V. I., Trigub, C., & Oram, S. (2013). Risk factors for mental disorders in women survivors of human trafficking: a historical cohort study. *BMC Psychiatry*, 13, 266. doi:10.1186/1471-244x-13-204
- Acarturk, C., Konuk, E., Cetinkaya, M., Senay, I., Sijbrandij, M., Gulen, B., & Cuijpers, P. (2016). The efficacy of eye movement desensitization and reprocessing for post-traumatic stress disorder and depression among Syrian refugees: Results of a randomized controlled trial. *Psychological Medicine*, 1-11. doi:10.1017/S0033291716001070
- Aggarwal, V. R., Macfarlane, G. J., Tajar, A., Mulvey, M. R., Power, A., Ray, D., & McBeth, J. (2014). Functioning of the hypothalamic-pituitary-adrenal and growth hormone axes in frequently unexplained disorders: Results of a population study. *European Journal of Pain*, 18, 447-454. doi:10.1002/j.1532-2149.2013.00413.x
- Anacker, C., O'Donnell, K. J., & Meaney, M. J. (2014). Early life adversity and the epigenetic programming of hypothalamic-pituitary-adrenal function. *Dialogues in Clinical Neuroscience*, 16, 321-333.
- Antonelli, M. C., Pallares, M. E., Ceccatelli, S., & Spulber, S. (2016). Long-term consequences of prenatal stress and neurotoxicants exposure on neurodevelopment. *Progress in Neurobiology*, doi:10.1016/j.pneurobio.2016.05.005
- Banovic, B., & Bjelajac, Z. (2012). Traumatic experiences, psychophysical consequences and needs of human trafficking victims. *Vojnosanitetski pregled*, 69, 94-97. <http://dx.doi.org/10.2298/VSP1201094B>
- Burke, N. N., Finn, D. P., McGuire, B. E., & Roche, M. (2016). Psychological stress in early life as a predisposing factor for the development of chronic pain: Clinical and preclinical evidence and neurobiological mechanisms. *Journal of Neuroscience Research*. doi:10.1002/jnr.23802
- Busso, D. S., McLaughlin, K. A., & Sheridan, M. A. (2016). Dimensions of adversity, physiological reactivity, and externalizing psychopathology in adolescence: Deprivation and threat. *Psychosomatic Medicine*. doi:10.1097/psy.0000000000000369
- Cannon, A. C., Arcara, J., Graham, L. M., & Macy, R. J. (2016). Trafficking and health: A systematic review of research methods. *Trauma Violence Abuse*. doi:10.1177/1524838016650187

- Cary, M., Oram, S., Howard, L. M., Trevillion, K., & Byford, S. (2016). Human trafficking and severe mental illness: An economic analysis of survivors' use of psychiatric services. *BMC Health Services Research*, 16, e157. doi:10.1186/s12913-016-1541-0
- Chesnay, M. (2013). Psychiatric-mental health nurses and the sex trafficking pandemic. *Issues in Mental Health Nursing*, 34, 901–907. doi:10.3109/01612840.2013.857200
- Chiapponi, C., Piras, F., Piras, F., Caltagirone, C., & Spalletta, G. (2016). GABA system in schizophrenia and mood disorders: A mini review on third-generation imaging studies. *Front Psychiatry*, 7, 61. doi:10.3389/fpsy.2016.00061
- Chung, R. J., & English, A. (2015). Commercial sexual exploitation and sex trafficking of adolescents. *Current Opinion in Pediatrics*, 27, 427–433. doi:10.1097/mop.0000000000000242
- Cohen, J. N. (2008). Using feminist, emotion-focused, and developmental approaches to enhance cognitive-behavioral therapies for posttraumatic stress disorder related to childhood sexual abuse. *Psychotherapy: Theory, Research, Practice, Training*, 45, 227–246. doi:10.1037/0033-3204.45.2.227
- Committee opinion No. 507: Human trafficking (2011). *Obstetrics & Gynecology*, 118, 767–770. doi:10.1097/AOG.0b013e3182310d06
- Corr, P. J., & Perkins, A. M. (2006). The role of theory in the psychophysiology of personality: From Ivan Pavlov to Jeffrey Gray. *International Journal of Psychophysiology*, 62, 367–376. doi:10.1016/j.ijpsycho.2006.01.005
- Crawford, M., & Kaufman, M. R. (2008). Sex trafficking in Nepal: Survivor characteristics and long-term outcomes. *Violence Against Women*, 14, 905–916. doi:10.1177/1077801208320906
- Cwikel, J., Chudakov, B., Paikin, M., Agmon, K., & Belmaker, R. H. (2004). Trafficked female sex workers awaiting deportation: comparison with brothel workers. *Archives of Women's Mental Health*, 7, 243–249. doi:10.1007/s00737-004-0062-8
- Davidson, L. L., Grigorenko, E. L., Boivin, M. J., Rapa, E., & Stein, A. (2015). A focus on adolescence to reduce neurological, mental health and substance-use disability. *Nature*, 527, S161–S166. doi:10.1038/nature16030
- Davis, J., Eyre, H., Jacka, F. N., Dodd, S., Dean, O., McEwen, S., ... Berk, M. (2016). A review of vulnerability and risks for schizophrenia: Beyond the two hit hypothesis. *Neuroscience & Biobehavioral Reviews*, 65, 185–194. doi:10.1016/j.neubiorev.2016.03.017
- Decker, M. R., Mack, K. P., Barrows, J. J., & Silverman, J. G. (2009). Sex trafficking, violence victimization, and condom use among prostituted women in Nicaragua. *International Journal of Gynecology & Obstetrics*, 107, 151–152. doi:10.1016/j.ijgo.2009.06.002
- del Rosario-Montejo, O., Molina-Rueda, F., Muñoz-Lasa, S., & Alguacil-Diego, I. M. (2015). Efectividad de la terapia ecuestre en niños con retraso psicomotor. *Neurología*, 30, 425–432. doi:10.1016/j.nrl.2013.12.023
- Demirdas, A., Naziroglu, M., & Unal, G. O. (2016). Agomelatine reduces brain, kidney and liver oxidative stress but increases plasma cytokine production in the rats with chronic mild stress-induced depression. *Metabolic Brain Disease*. doi:10.1007/s11011-016-9874-2
- Diaz Heijtz, R., & Forsberg, H. (2015). Translational studies exploring neuroplasticity associated with motor skill learning and the regulatory role of the dopamine system. *Developmental Medicine & Child Neurology*, 57, 10–14. doi:10.1111/dmnc.12692
- Diehl, A., Pillon, S. C., dos Santos, M. A., Rassool, G. H., & Laranjeira, R. (2016). Criminality and sexual behaviours in substance dependents seeking treatment. *Journal of Psychoactive Drugs*, 48, 124–134. doi:10.1080/02791072.2016.1168534
- Ehlert, U. (2013). Enduring psychobiological effects of childhood adversity. *Psychoneuroendocrinology*, 38, 1850–1857. doi:10.1016/j.psyneuen.2013.06.007
- English, A. (2015). Human trafficking of children and adolescents. *JAMA Pediatrics*, 169, e152283. doi:10.1001/jamapediatrics.2015.2283
- Ford, J. D., & Kidd, P. (1998). Early childhood trauma and disorders of extreme stress as predictors of treatment outcome with chronic posttraumatic stress disorder. *Journal of Traumatic Stress*, 11, 743–761. doi:10.1023/a:1024497400891
- Frances, A., Pincus, H. A., Widiger, T. A., Davis, W. W., & First, M. B. (1990). DSM-IV: Work in progress. *American Journal of Psychiatry*, 147, 1439–1448. doi:10.1176/ajp.147.11.1439
- Froehlich, M. A. (1984). A comparison of the effect of music therapy and medical play therapy on the verbalization behavior of pediatric patients. *Journal of Music Therapy*, 21, 2–15. <http://dx.doi.org/10.1093/jmt/21.1.2>
- Gibbons, P., & Stoklosa, H. (2016). Identification and treatment of human trafficking victims in the emergency department: A case report. *The Journal of Emergency Medicine*, 50, 715–719. doi:10.1016/j.jemermed.2016.01.004
- Grace, A. M., Lippert, S., Collins, K., Pineda, N., Tolani, A., Walker, R., ... Horwitz, S. M. (2014). Educating health care professionals on human trafficking. *Pediatric Emergency Care*, 30, 856–861. doi:10.1097/pec.0000000000000287
- Greenbaum, V. J., Dodd, M., & McCracken, C. (2015). A short screening tool to identify victims of child sex trafficking in the health care setting. *Pediatric Emergency Care*. doi:10.1097/pec.0000000000000602
- Gupta, K. K., Gupta, V. K., & Shirasaka, T. (2016). An update on fetal alcohol syndrome-pathogenesis, risks, and treatment. *Alcoholism Clinical and Experimental Research*. doi:10.1111/acer.13135
- Helton, M. (2016). Human trafficking: How a joint task force between health care providers and law enforcement can assist with identifying victims and prosecuting traffickers. *Health Matrix Cleveland*, 26, 433–473.
- Hewagalamlage, S. D., Clarke, I. J., Rao, A., & Henry, B. A. (2016). Ewes with divergent Cortisol responses to Adrenocorticotropin (ACTH) exhibit functional differences in the Hypothalamo-Pituitary-Adrenal (HPA) axis. *Endocrinology*, en20161287. doi:10.1210/en.2016-1287
- Hossain, M., Zimmermann, C., Abas, M., Light, M., & Watts, C. (2010). The relationship of trauma to mental disorders among trafficked and sexually exploited girls and women. *American Journal of Public Health*, 100, 2442–2449. doi:10.2105/ajph.2009.173229
- Houston-Kolnik, J. D., Todd, N. R., & Wilson, M. (2016). Preliminary validation of the sex trafficking attitudes scale. *Violence Against Women*. doi:10.1177/1077801215621178
- Huston, H. R., Anglin, D., & Eckstein, M. (1996). Drive-by shootings by violent street gangs in Los Angeles: a five-year review from 1989 to 1993. *Academic Emergency Medicine*, 3, 300–303.
- Hylland, P., Shevlin, M., Elklit, A., Murphy, J., Vallieres, F., Garvert, D. W., & Cloitre, M. (2016). An assessment of the construct validity of the ICD-11 proposal for complex posttraumatic stress disorder. *Psychological Traum*. doi:10.1037/tra0000114
- Ijadi-Maghsoudi, R., Cook, M., Barnert, E. S., Gaboian, S., & Bath, E. (2016). Understanding and responding to the needs of commercially sexually exploited youth. *Child and Adolescent Psychiatric Clinics of North America*, 25, 107–122. doi:10.1016/j.chc.2015.08.007
- ILO Global Estimate of Forced Labour: Results and methodology/International Labour Office, Special Action Programme to Combat Forced Labour (SAP-FL). (2012). Geneva: ILO 2012. Retrieved from <http://apflnet.ilo.org/resources/ilo-global-estimate-of-forced-labour-2012-results-and-methodology>

- Institute of Medicine & National Research Council. (2014). *Confronting commercial sexual exploitation and sex trafficking of minors in the United States: A guide for the legal sector*. Washington, DC: National Academies Press. Copyright 2014 by the National Academy of Sciences. All rights reserved.
- Jana, S., Dey, B., Reza-Paul, S., & Steen, R. (2014). Combating human trafficking in the sex trade: can sex workers do it better? *Journal of Public Health*, 36, 622–628. doi:10.1093/pubmed/fdt095
- Kangaslampi, S., Garoff, F., & Peltonen, K. (2015). Narrative exposure therapy for immigrant children traumatized by war: study protocol for a randomized controlled trial of effectiveness and mechanisms of change. *BMC Psychiatry*, 15, 553. doi:10.1186/s12888-015-0520-z
- Karatzias, T., Shevlin, M., Fyvie, C., Hyland, P., Efthymiadou, E., Wilson, D., ... Cloitre, M. (2016). An initial psychometric assessment of an ICD-11 based measure of PTSD and complex PTSD (ICD-TQ): Evidence of construct validity. *Journal of Anxiety Disorders*, 44, 73–79. doi:10.1016/j.janxdis.2016.10.009
- Keeshin, B. R., & Strawn, J. R. (2014). Psychological and pharmacologic treatment of youth with posttraumatic stress disorder. *Child and Adolescent Psychiatric Clinics of North America*, 23, 399–411. doi:10.1016/j.chc.2013.12.002
- Kimerling, R., Alvarez, J., Pavao, J., Kaminski, A., & Baumrind, N. (2007). Epidemiology and consequences of women's revictimization. *Women's Health Issues*, 17, 101–106. <http://dx.doi.org/10.1016/j.whi.2006.12.002>
- Kliner, M., & Stroud, L. (2012). Psychological and health impact of working with victims of sex trafficking. *Journal of Occupational Health*, 54, 9–15. <http://dx.doi.org/10.1539/joh.11-0125-OA>
- Kumar, J. R., Rajkumar, R., Lee, L. C., & Dawe, G. S. (2016). Nucleus incertus contributes to an anxiogenic effect of buspirone in rats: Involvement of 5-HT1A receptors. *Neuropharmacology*. doi:10.1016/j.neuropharm.2016.07.019
- Larsen, S. E., Wiltsey Stirman, S., Smith, B. N., & Resick, P. A. (2016). Symptom exacerbations in trauma-focused treatments: Associations with treatment outcome and non-completion. *Behaviour Research and Therapy*, 77, 68–77. doi:10.1016/j.brat.2015.12.009
- Lebois, L. A., Wolff, J. D., & Ressler, K. J. (2016). Neuroimaging genetic approaches to posttraumatic stress disorder. *Experimental Neurology*. doi:10.1016/j.expneurol.2016.04.019
- Li-Tempel, T., Larra, M. F., Winnikes, U., Tempel, T., DeRijk, R. H., Schulz, A., ... Schote, A. B. (2016). Polymorphisms of genes related to the hypothalamic-pituitary-adrenal axis influence the cortisol awakening response as well as self-perceived stress. *Biological Psychology*. doi:10.1016/j.biopsycho.2016.07.010
- Liu, G., Zhu, T., Zhang, A., Li, F., Qian, W., & Qian, B. (2016). Heightened stress response and cognitive impairment after repeated neonatal sevoflurane exposures might be linked to excessive GABAAR-mediated depolarization. *Journal of Anesthesia*. doi:10.1007/s00540-016-2215-0
- Lutz, P. E., Almeida, D., Fiori, L. M., & Turecki, G. (2015). Childhood maltreatment and stress-related psychopathology: The epigenetic memory hypothesis. *Current Pharmaceutical Design*, 21, 1413–1417. <http://dx.doi.org/10.2174/1381612821666150105124928>
- Mantini, L. (2008). Human trafficking of Amerindian women in Guyana: challenges and strategies. *International Nursing Review*, 55, 341–348. doi:10.1111/j.1466-7657.2008.00624.x
- Marina, D., Klose, M., Nordenbo, A., Liebach, A., & Feldt-Rasmussen, U. (2015). Early endocrine alterations reflect prolonged stress and relate to 1-year functional outcome in patients with severe brain injury. *European Journal of Endocrinology*, 172, 813–822. doi:10.1530/eje-14-1152
- Matošić, A., Marusic, S., Vidrih, B., Kovak-Mufic, A., & Cicin-Sain, L. (2016). Neurobiological bases of alcohol addiction. *Acta Clinica Croatica*, 55, 134–150. <http://dx.doi.org/10.20471/acc.2016.55.01.19>
- McEwen, B. S., Gray, J. D., & Nasca, C. (2015). 60 years of neuroendocrinology: Redefining neuroendocrinology: stress, sex and cognitive and emotional regulation. *Journal of Endocrinology*, 226, T67–T83. doi:10.1530/joe-15-0121
- McEwen, B. S., & Wingfield, J. C. (2003). The concept of allostasis in biology and biomedicine. *Hormones and Behavior*, 43, 2–15. [http://dx.doi.org/10.1016/S0018-506X\(02\)00024-7](http://dx.doi.org/10.1016/S0018-506X(02)00024-7)
- Mills, B., Carter, O., Rudd, C., Claxton, L., & O'Brien, R. (2016). An experimental investigation into the extent social evaluation anxiety impairs performance in simulation-based learning environments amongst final-year undergraduate nursing students. *Nurse Education Today*, 45, 9–15. doi:10.1016/j.nedt.2016.06.006
- Minnesota Department of Health. (2014). *Safe Harbor Minnesota*. Retrieved from <http://www.health.state.mn.us/injury/topic/safeharbor/>
- Mogami, S., Sadakane, C., Nahata, M., Mizuhara, Y., Yamada, C., Hattori, T., & Takeda, H. (2016). CRF receptor 1 antagonism and brain distribution of active components contribute to the ameliorative effect of rikkunshito on stress-induced anorexia. *Scientific Reports*, 6, 27516. doi:10.1038/srep27516
- Nijdam, M. J., & Olff, M. (2016). Erasing memory traces of trauma with eye movement desensitization and reprocessing therapy. *European Journal of Psychotraumatology*, 7, 32545. doi:10.3402/ejpt.v7.32545
- Nuzzolese, E. (2014). Human trafficking: Role of oral health care providers. *The Journal of Forensic Odontostomatology*, 32, 1–8.
- Oram, S., Ostrovschi, N. V., Gorceag, V. I., Hotineanu, M. A., Gorceag, L., Trigub, C., & Abas, M. (2012). Physical health symptoms reported by trafficked women receiving post-trafficking support in Moldova: prevalence, severity and associated factors. *BMC Women's Health*, 12, 266. doi:10.1186/1472-6874-12-20
- Oram, S., Stöckl, H., Busza, J., Howard, L. M., & Zimmerman, C. (2012). Prevalence and risk of violence and the physical, mental, and sexual health problems associated with human trafficking: Systematic review. *PLoS Medicine*, 9, e1001224. doi:10.1371/journal.pmed.1001224
- Osterlund, C. D., Rodriguez-Santiago, M., Woodruff, E. R., Newsom, R. J., Chadayammuri, A. P., & Spencer, R. L. (2016). Glucocorticoid fast feedback inhibition of stress-induced ACTH secretion in the male rat: Rate independence and stress-state resistance. *Endocrinology*, 157, 2785–2798. doi:10.1210/en.2016-1123
- Ostrovski, N. V., Prince, M. J., Zimmerman, C., Hotineanu, M. A., Gorceag, L. T., Gorceag, V. I., ... Abas, M. A. (2011). Women in post-trafficking services in moldova: Diagnostic interviews over two time periods to assess returning women's mental health. *BMC Public Health*, 11, 564. doi:10.1186/1471-2458-11-232
- Provençal, N., & Binder, E. B. (2015). The effects of early life stress on the epigenome: From the womb to adulthood and even before. *Experimental Neurology*, 268, 10–20. doi:10.1016/j.expneurol.2014.09.001
- Provenzi, L., Giusti, L., Fumagalli, M., Tasca, H., Ciceri, F., Menozzi, G., ... Montiroso, R. (2016). Pain-related stress in the Neonatal Intensive Care Unit and salivary cortisol reactivity to socio-emotional stress in 3-month-old very preterm infants. *Psychoneuroendocrinology*, 72, 161–165. doi:10.1016/j.psyneuen.2016.07.010
- Ratzer, M., Brink, O., Knudsen, L., & Elklit, A. (2014). Posttraumatic stress in intensive care unit survivors - a prospective study. *Health Psychology and Behavioral Medicine*, 2, 882–898. doi:10.1080/21642850.2014.943760

- Recommendations to reduce psychological harm from traumatic events among children and adolescents. (2008). *American Journal of Preventive Medicine*, 35, 314–316. doi:10.1016/j.amepre.2008.06.025
- Richards, T. A. (2014). Health implications of human trafficking. *Nursing for Women's Health*, 18, 155–162. doi:10.1111/1751-486x.12112
- Rimal, R., & Papadopoulos, C. (2016). The mental health of sexually trafficked female survivors in Nepal. *International Journal of Social Psychiatry*. doi:10.1177/0020764016651457
- Roe-Sepowitz, D. E., Gallagher, J., Risinger, M., & Hickle, K. (2015). The sexual exploitation of girls in the United States: The role of female pimps. *Journal of Interpersonal Violence*, 30, 2814–2830. doi:10.1177/0886260514554292
- Ross, C., Dimitrova, S., Howard, L. M., Dewey, M., Zimmerman, C., & Oram, S. (2015). Human trafficking and health: A cross-sectional survey of NHS professionals' contact with victims of human trafficking. *BMJ Open*, 5, e008682. doi:10.1136/bmjopen-2015-008682
- Sachser, C., Keller, F., & Goldbeck, L. (2016). Complex PTSD as proposed for ICD-11: Validation of a new disorder in children and adolescents and their response to Trauma-Focused Cognitive Behavioral Therapy. *Journal of Child Psychology and Psychiatry*. doi:10.1111/jcpp.12640
- Savendahl, L. (2012). The effect of acute and chronic stress on growth. *Science Signaling*, 5(247), pt9. doi:10.1126/scisignal.2003484
- Schaefer, S. E., Camacho-Gomez, R., Sadeghi, B., Kaiser, L., German, J. B., & de la Torre, A. (2015). Assessing Child Obesity and Physical Activity in a Hard-to-Reach Population in California's Central Valley, 2012–2013. *Preventing Chronic Disease*, 12, E117. doi:10.5888/pcd12.140577
- Serafini, G., Muzio, C., Piccinini, G., Flouri, E., Ferrigno, G., Pompili, M., ... Amore, M. (2015). Life adversities and suicidal behavior in young individuals: a systematic review. *European Child & Adolescent Psychiatry*, 24, 1423–1446. doi:10.1007/s00787-015-0760-y
- Shandro, J., Chisolm-Straker, M., Duber, H. C., Findlay, S. L., Munoz, J., Schmitz, G., ... Wingkun, N. (2016). Human trafficking: A guide to identification and approach for the emergency physician. *Annals of Emergency Medicine*. doi:10.1016/j.annemergmed.2016.03.049
- Simchon Tenenbaum, Y., Weizman, A., & Rehavi, M. (2015). The impact of chronic early administration of psychostimulants on brain expression of BDNF and other neuroplasticity-relevant proteins. *Journal of Molecular Neuroscienc.* doi:10.1007/s12031-015-0611-9
- Szklarczyk, K., Korostynski, M., Golda, S., Piechota, M., Ficek, J., & Przewlocki, R. (2016). Endogenous opioids regulate glucocorticoid-dependent stress-coping strategies in mice. *Neuroscience*, 330, 121–137. doi:10.1016/j.neuroscience.2016.05.034
- Thakare, V. N., Dhakane, V. D., & Patel, B. M. (2016). Potential antidepressant-like activity of silymarin in the acute restraint stress in mice: Modulation of corticosterone and oxidative stress response in cerebral cortex and hippocampus. *Pharmacological Reports*, 68, 1020–1027. doi:10.1016/j.pharep.2016.06.002
- Tsutsumi, A., Izutsu, T., Poudyal, A. K., Kato, S., & Marui, E. (2008). Mental health of female survivors of human trafficking in Nepal. *Social Science & Medicine*, 66, 1841–1847. doi:10.1016/j.socscimed.2007.12.025
- United Nations Office on Drugs and Crime [UNODC]. (2009). *Global report on trafficked persons*. Vienna. Retrieved from https://www.unodc.org/documents/wdr/WDR_2009/WDR2009_eng_web.pdf
- US Congress. (2014). *H.R.4980 - Preventing sex trafficking and strengthening families act*. Public Law No: 113–183. C.F.R.
- van der Kolk, B. A. (2001). The assessment and treatment of complex PTSD. In R. Yehuda (Ed.), *Traumatic stress* (pp. 1–29). Arlington, VA: American Psychiatric Press.
- Vyas, S., Rodrigues, A. J., Silva, J. M., Tronche, F., Almeida, O. F., Sousa, N., & Sotiropoulos, I. (2016). Chronic stress and glucocorticoids: From neuronal plasticity to neurodegeneration. *Neural Plasticity*, 2016, 6391686. doi:10.1155/2016/6391686
- Walter, K. H., Buckley, A., Simpson, J. M., & Chard, K. M. (2014). Residential PTSD treatment for female veterans with military sexual trauma: Does a history of childhood sexual abuse influence outcome? *Journal of Interpersonal Violence*, 29, 971–986. doi:10.1177/0886260513506055
- Wilson, M. A., Grillo, C. A., Fadel, J. R., & Reagan, L. P. (2015). Stress as a one-armed bandit: Differential effects of stress paradigms on the morphology, neurochemistry and behavior in the rodent amygdala. *Neurobiology of Stress*, 1, 195–208. doi:10.1016/j.yjnstr.2015.06.001
- Windholz, G. (1987). Pavlov as a psychologist. A reappraisal. *Pavlovian Journal of Biological Science*, 22, 103–112.
- Wong, M. (2008). Interventions to reduce psychological harm from traumatic events among children and adolescents. *American Journal of Preventive Medicine*, 35, 398–400. doi:10.1016/j.amepre.2008.07.002
- Wood, C. S., Valentino, R. J., & Wood, S. K. (2016). Individual differences in the locus coeruleus-norepinephrine system: relevance to stress-induced cardiovascular vulnerability. *Physiology & Behavior*. doi:10.1016/j.physbeh.2016.07.008
- Xia, L., & Yao, S. (2015). The involvement of genes in adolescent depression: A systematic review. *Frontiers in Behavioral Neuroscience*, 9, 329. doi:10.3389/fnbeh.2015.00329
- Yang, L., Xu, T., Zhang, K., Wei, Z., Li, X., Huang, M., ... Cai, X. (2016). The essential role of hippocampal alpha6 subunit-containing GABAA receptors in maternal separation stress-induced adolescent depressive behaviors. *Behavioural Brain Research*. doi:10.1016/j.bbr.2016.07.002
- Yu, C., & McClellan, J. (2016). Genetics of substance use disorders. *Child and Adolescent Psychiatric Clinics of North America*, 25, 377–385. doi:10.1016/j.chc.2016.02.002
- Zimmerman, C., Hossain, M., & Watts, C. (2011). Human trafficking and health: A conceptual model to inform policy, intervention and research. *Social Science & Medicine*, 73, 327–335. doi:10.1016/j.socscimed.2011.05.028
- Zimmerman, C., Hossain, M., Yun, K., Gajdzdziej, V., Guzun, N., Tchomarova, M., ... Watts, C. (2008). The health of trafficked women: A survey of women entering posttrafficking services in Europe. *American Journal of Public Health*, 98, 55–59. doi:10.2105/ajph.2006.108357



© 2017 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



Cogent Medicine (ISSN: 2331-205X) is published by Cogent OA, part of Taylor & Francis Group.

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

